

059 Keishibukuryogankayokuinin

| | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | |
|---|--|---|---|----------------------------|--|--|----------------|-----------------------------------|-------------|---|--|-----------------|------------------|-------------|--------------|------------|----------|
| | | | | | | | | | | | | Cinnamon Bark | Poria Sclerotium | Moutan Bark | Peach Kernel | Peony Root | Cox Seed |
| 1 | Revised: May 2007 (4th version) | TSUMURA Keishibukuryogankayokuinin Extract Granules for Ethical Use Tsumura & Co. | (62AM)578 Oct 1987 Oct 1987 | 7.5 | 3.75 | JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate | Granules | TSUMURA/125 | A | before or between meals, 2 or 3 times | 500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets | 4.0 | 4.0 | 4.0 | 4.0 | 4.0 | 10.0 |

Table of Indications

| | |
|---|---|
| A | The following symptoms of those patients with a comparatively strong constitution who sometimes complain of lower abdominal pain, shoulder stiffness, dull headache, dizziness, cold feet with feeling of hot flush, etc.: Menstrual irregularity, automatic imbalance syndrome peculiar to women resembling climacteric disturbance, acne, spots, and roughness of the hands and feet |
|---|---|