

SEISHINRENSHIIN (清心蓮子飲)

Indications

Following symptoms in persons with moderate or poor physical strength who have gastrointestinal weakness, general malaise, dry mouth or tongue and difficulty urinating:

Frequent urination, feeling of residual urine, urination pain, urination difficulty, cloudy urine, and leukorrhea.

! Caution for use

To be avoided

Don't give this medicine to children under 3 months of age.

Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - Persons undergoing medical treatment from a physician.
 - Pregnant women or women suspected of being pregnant.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Symptoms' name	Symptoms
Interstitial pneumonia	Shortness and / or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance
Hepatic function failure	The following symptoms appear: Fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite

- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.

Components

Seishinrenshiin (清心蓮子飲) is extract from mixture of the following crude drugs;

Nelumbo Seed (蓮肉), Ophiopogon Root (麥門冬), Poria Sclerotium (茯苓), Ginseng (人參), Plantago Seed (車前子), Scutellaria Root (黃芩), Astragalus Root (黃耆), Lycium Bark (地骨皮), Glycyrrhiza (甘草).

*If you have any questions, please consult to a physician, pharmacist, or registered distributor .
Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University