

003 Bofutsushosan

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio																		
												Japanese Angelica Root	Peony Root	Chidum Rhizome	Gardenia Fruit	Forsythia Fruit	Menhadi Herb	Ginger	Fresh Ginger	Schizonepeta Spike	Rhizome	Saposhnikovia Root and Rhizome	Glehnia Root and Rhizome	Epilobium Herb	Rhubarb	Sodium Sulfate Hydrate	Anhydrous Sodium Sulfate	Anacardolides Rhizome	Plyosodon Root	Scutellaria Root
1	Revised: Feb 2018 (9th version)	Honzo Bofutsushosan Extract Granules-M Honzo Pharmaceutical Co., Ltd.	(61AM)4158 Oct 1986 Oct 1986	7.5	5.0	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate	Granules	H62	A	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	1.2	1.2	1.2	1.2	1.2	1.2	0.4	1.2	1.2	1.2	1.5	1.5	2.0	2.0	2.0	2.0	2.0	2.0	3.0
2	Revised: Feb 2018 (9th version)	JPS Bofutsushosanryo Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd.	62AM-0094 Oct 1987 Oct 1987	7.5	5.0	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-62)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	1.2	1.2	1.2	1.2	1.2	1.2	0.3	1.2	1.2	1.2	1.5	1.5	2.0	2.0	2.0	2.0	2.0	2.0	3.0
3	Revised: Feb 2018 (11th version)	Kotaro Bofutsushosan Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3960 Oct 1986 Oct 1986	9.0	6.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N62	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	1.2	1.2	1.2	1.2	1.2	1.2	0.3	1.2	1.2	1.2	1.5	0.7	2.0	2.0	2.0	2.0	2.0	2.0	3.0
4	Revised: Feb 2018 (5th version)	Kracie Bofutsushosanryo Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3574 Jul 2007 Jul 2007	7.5	5.7	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-62 3.75g EK-62 2.5g	A	before or between meals, 2 or 3 times	3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g	1.2	1.2	1.2	1.2	1.2	1.2	0.4	1.2	1.2	1.2	1.5	0.75	2.0	2.0	2.0	2.0	2.0	2.0	3.0
5	Revised: Feb 2018 (5th version)	Kracie Bofutsushosan Extract Tablets Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00632000 Jul 2007 Jul 2007	8.91 (27 tablets)	5.5	JP Magnesium Stearate, JP Carmellose Calcium, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose	Tablets	(EKT-62)	A	before or between meals, 2 or 3 times	378 tablets (3 tablets x 6 packets x 21 sheets) 1,764 tablets (3 tablets x 6 packets x 98 sheets)	1.2	1.2	1.2	1.2	1.2	1.2	0.4	1.2	1.2	1.2	1.5	0.75	2.0	2.0	2.0	2.0	2.0	2.0	3.0
6	Revised: Feb 2018 (6th version)	Matsuura Bofutsushosan Extract Granules Matsuura Yakugyo Co., Ltd.	(61AM)4177 Oct 1987 Oct 1987	7.5	viscous extract 10.0*	Hypromellose, Macrogol 6000, Lactose Hydrate, Dextrin, Corn Starch	Granules	62, Matsuura	A	before or between meals, 2 or 3 times	500g 2.5g x 84 packets 2.5g x 300 packets	1.2	1.2	1.2	1.2	1.2	1.2	0.4	1.2	1.2	1.2	1.5	1.5	2.0	2.0	2.0	2.0	2.0	2.0	3.0
7	Revised: Feb 2018 (2nd version)	OHSUGI Bofutsushosan Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3890 Oct 1986 Oct 1986	9.0	5.2	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-62	A	before or between meals, 2 or 3 times	500g 882g (3.0g x 294 packets) 252g (3.0g x 84 packets)	1.2	1.2	1.2	1.2	1.2	1.2	0.3	1.2	1.2	1.2	1.5	0.7	2.0	2.0	2.0	2.0	2.0	2.0	3.0
8	Revised: Feb 2018 (7th version)	SANWA Bofutsushosanryo Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3628 Oct 1986 Nov 1986	9.0	5.4	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(S-26)	C	before or between meals, 3 times	500g 3g x 300 packets	1.2	1.2	1.2	1.2	1.2	1.2	0.3	1.2	1.2	1.2	1.5	0.75	2.0	2.0	2.0	2.0	2.0	2.0	3.0
9	Revised: Feb 2018 (7th version)	Taikodo Bofutsushosanryo Extract Granules Taikoseido Pharmaceutical Co., Ltd.	(61AM)3918 Oct 1986 Nov 1986	7.5	5.4	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-62	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.5	1.5	2.0	2.0	2.0	2.0	2.0	2.0	3.0
10	Revised: Feb 2018 (11th version)	TEIKOKU Bofutsushosan Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3696 Oct 1987 Oct 1987	7.5	3.04	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 62	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	1.2	1.2	1.2	1.2	1.2	1.2	0.4	1.2	1.2	1.2	1.5	1.5	2.0	2.0	2.0	2.0	2.0	2.0	3.0
11	Revised: Feb 2018 (9th version)	Toyo Bofutsushosanryo Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AMY)0391 Oct 1987 Apr 1988	7.5	5.0	Corn Starch	Fine granules	TY-100	A	on empty stomach, 3 times	500g 2.5g x 252 packets	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.5	1.5	2.0	2.0	2.0	2.0	2.0	2.0	3.0
12	Revised: Feb 2018 (9th version)	TSUMURA Bofutsushosan Extract Granules for Ethical Use Tsumura & Co.	(61AM)1172 Oct 1986 Oct 1986	7.5	4.5	JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/62	A	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	1.2	1.2	1.2	1.2	1.2	1.2	0.3	1.2	1.2	1.2	1.5	0.7	2.0	2.0	2.0	2.0	2.0	2.0	3.0

*: correspond to 5.0g of dry extract

Table of Indications

A	The following symptoms of those patients with thick subcutaneous fat in the abdomen and a tendency to constipation: Accessory symptoms associated with hypertension (palpitation, shoulder stiffness, and hot flushes), obesity, swelling, and constipation
B	The following symptoms of those patients with thick subcutaneous fat, constipation, and decreased urinary output: Chronic constipation, hyperchlorhydria, kidney disease, cardiac debility, arteriosclerosis, hypertension, cerebral hemorrhage, and shoulder stiffness due to these diseases
C	The following symptoms of those patients with thick subcutaneous fat who have constipation, or pyrosis, shoulder stiffness, decreased urine volume, etc.: Obesity, hypertension, chronic constipation, hemorrhoids, chronic nephritis, and eczema