

005 Bukuryoin

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio						
												Poria Sclerotium	Atractylodes Rhizome	Atractylodes Lancea	Ginseng	Ginger	Citrus Urshiu Peel	Immature Orange
1	Revised: Feb 2008 (5th version)	<b>Kotaro Bukuryoin Extract Fine Granules</b>  Kotaro Pharmaceutical Co., Ltd.	(61AM)3837 Oct 1986 Oct 1986	6.0	3.8	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N69	A	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	5.0	4.0	/	3.0	0.8	3.0	1.5
2	Revised: Mar 2013 (5th version)	<b>TSUMURA Bukuryoin Extract Granules for Ethical Use</b>  Tsumura & Co.	(61AM)1130 Oct 1986 Oct 1986	7.5	2.75	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/69	B	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	5.0	/	4.0	3.0	1.0	3.0	1.5

**Table of Indications**

A	The following symptoms of those patients with blocked feeling in the stomach, stomach fullness, excessive gastric fluid, nausea, vomiting, anorexia, and decreased urinary output: Gastritis, gastropnoia, gastric atony, gastric neurosis, gastric dilatation, esophageal achalasia, and dyspepsia
B	The following symptoms of those patients with nausea or heartburn and decreased urine volume: Gastritis, gastric atony, and excessive fluid retention in the stomach