

007 Bushirichuto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio				
												Ginseng	Processed Aconite Root	Processed Ginger	Glycyrrhiza	Atractylodes Rhizome
1	Revised: Jun 2016 (4th version)	SANWA Bushirichuto Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3614 Oct 1986 Nov 1986	4.5	2.8	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(S-09)	A	before or between meals, 3 times	500g 1.5g x 300 packets	3.0	1.0	3.0	3.0	3.0
2	Revised: Jul 2010 (5th version)	SANWA Bushirichuto Extract Fine Granules Sanwa Shoyaku Co., Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3614 Oct 1986 Nov 1986	4.5	2.8	JP Lactose Hydrate, JP Corn Starch, JP Microcrystalline Cellulose, JP Light Anhydrous Silicic Acid, Partly Pregelatinized Starch	Fine granules	(EK-410)	A	before or between meals, 3 times	500g 1.5g x 42 packets 1.5g x 294 packets	3.0	1.0	3.0	3.0	3.0

Table of Indications

A	The following symptoms of those patients with gastrointestinal weakness, a bad complexion, a face without vitality, high urine output, cold feeling in the extremities, tendency to diarrhea, frequent nausea, dizziness, dull headache and stomachache: Chronic gastrointestinal catarrh and gastric atony
---	--