

011 Choreito

| | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | |
|----|--|--|--|----------------------|-------------------------------------|--|---------------|-----------------------------|-------------|---|--|----------------------|------------------|--|--------------|-------------|---------|
| | | | | | | | | | | | | Polyporus Sclerotium | Poria Sclerotium | Aluminum Silicate Hydrate with Silicon Dioxide | Aislma Tuber | Donkey Glue | Gelatin |
| 1 | Revised: Jul 2011 (4th version) | Honzo Choreito Extract Granules-M Honzo Pharmaceutical Co., Ltd. | (61AM)4818 Oct 1987 Oct 1987 | 7.5 | 2.4 | Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate | Granules | H40 | A | before or between meals, 3 times | 500g 2.5g x 294 packets 2.5g x 42 packets | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | / |
| 2 | Revised: Sep 2016 (5th version) | JPS Choreito Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd. | 61AM-4660 Oct 1987 Oct 1987 | 7.5 | 5.2 | Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate | Granules | (J-40) | A | before or between meals, 2 or 3 times | 500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets) | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | / |
| 3 | Revised: Jun 2012 (6th version) | Kotaro Choreito Extract Fine Granules Kotaro Pharmaceutical Co., Ltd. | (61AM)4662 Oct 1987 Oct 1987 | 6.0 | 1.2* | Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate | Fine granules | N40 | B | before or between meals, 2 or 3 times | 500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g) | 3.0 | 3.0 | 3.0 | 3.0 | / | 3.0 |
| 4 | Revised: Jan 2012 (3rd version) | Kracie Choreito Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd. | (61AM)4214 Jul 2007 Jul 2007 | 6.0 | 2.5 | JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide | Fine granules | KB-40 3.0g EK-40 2.0g | A | before or between meals, 2 or 3 times | 3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | / |
| 5 | Revised: Apr 2016 (3rd version) | Matsuura Choreito Extract Granules Matsuura Yakugyo Co., Ltd. | (63AM)406 Jul 1988 Jul 1988 | 6.0 | 0.7* | Hypromellose, Magnesium Stearate, Dextrin, Lactose Hydrate | Granules | 40, Matsuura | A | before or between meals, 2 or 3 times | 500g 2g x 84 packets 2g x 300 packets | 3.0 | 3.0 | 3.0 | 3.0 | / | 3.0 |
| 6 | Oct 2005 (1st version) | OHSUGI Choreito Extract Granules G Ohsugi Pharmaceutical Co., Ltd. | (62AM)0325 Oct 1987 Oct 1987 | 6.0 | 0.4* | Lactose Hydrate, Corn Starch, Magnesium Stearate | Granules | SG-40 | A | before or between meals, 2 or 3 times | 500g 588g (2.0g x 294 packets) 168g (2.0g x 84 packets) | 3.0 | 3.0 | 3.0 | 3.0 | / | / |
| 7 | Revised: Feb 2014 (4th version) | SANWA Choreito Extract Fine Granules Sanwa Shoyaku Co., Ltd. | (61AM)3631 Oct 1986 Nov 1986 | 7.5 | 3.7 | Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid | Fine granules | (S-34) | C | before or between meals, 3 times | 500g 2.5g x 300 packets | 3.0 | 3.0 | 3.0 | 3.0 | / | 3.0 |
| 8 | Revised: Jun 2011 (4th version) | Taikodo Choreito Extract Granules Taikoseido Pharmaceutical Co., Ltd. | (62AM)0178 Oct 1987 Oct 1987 | 6.0 | 4.2 | JP Lactose Hydrate, JP Magnesium Stearate | Granules | TM-40 | A | before or between meals, 3 times | 500g 2.0g x 84 packets 2.0g x 315 packets | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | / |
| 9 | Revised: Aug 2017 (7th version) | TEIKOKU Choreito Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd. | (62AM)0068 Oct 1987 Oct 1987 | 7.5 | 2.49 | Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate | Granules | TEIKOKU 40 | A | before meals, 3 times | 500g 2.5g x 42 packets 2.5g x 252 packets | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | / |
| 10 | Revised: Oct 2005 (4th version) | Toyo Choreito Extract Fine Granules Toyo Yakuko Co., Ltd. | (62AMY)0395 Jul 1988 Oct 1988 | 6.0 | 4.0 | Corn Starch | Fine granules | TY-080 | A | on empty stomach, 3 times | 500g 2.0g x 252 packets | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | / |
| 11 | Revised: Mar 2013 (5th version) | TSUMURA Choreito Extract Granules for Ethical Use Tsumura & Co. | (61AM)3282 Oct 1986 Oct 1986 | 7.5 | 2.5 | JP Magnesium Stearate, JP Lactose Hydrate | Granules | TSUMURA/40 | D | before or between meals, 2 or 3 times | 500g 2.5g x 42 packets 2.5g x 189 packets | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | / |

*: extract weight without Donkey Glue or Gelatin

Table of Indications

| | |
|---|---|
| A | The relief of decreased urine volume, difficulty in micturition, and micturition pain or feeling of residual urine |
| B | The following symptoms of those patients with thirst, painful micturition, dysuria, red urine or blood in urine, edema of lower back or legs: Nephritis, nephrosis, bladder catarrh, urethritis, and dysuria due to renal/bladder calculus |
| C | Cystitis, especially of acute cystitis, nephritis, dry mouth due to renal calculus or urethritis, increased urinary frequency, and micturition pain |
| D | The following symptoms of those patients who complain of decreased urine volume, dysuria, and oral dryness: Urethritis, nephritis, renal calculus, hemorrhagic inflammation, micturition pain, hematuria, edema of the lower part of the body from the waist down, feeling of residual urine, and diarrhea |