

011 Choreito

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio				
												Polyporus Sclerotium	Poria Sclerotium	Aluminum Silicate Hydrate with Silicon Dioxide	Alisma Tuber	Donkey Glue
1	Revised: Jul 2011 (4th version)	<b>Honzo Choreito Extract Granules-M</b> Honzo Pharmaceutical Co., Ltd.	(61AM)4818 Oct 1987 Oct 1987	7.5	2.4	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate	Granules	H40	A	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	3.0	3.0	3.0	3.0	3.0
2	Revised: Sep 2016 (5th version)	<b>JPS Choreito Extract Granules for Ethical Use</b> JPS Pharmaceutical Co., Ltd.	61AM-4660 Oct 1987 Oct 1987	7.5	5.2	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-40)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	3.0	3.0	3.0	3.0	3.0
3	Revised: Jun 2012 (6th version)	<b>Kotaro Choreito Extract Fine Granules</b> Kotaro Pharmaceutical Co., Ltd.	(61AM)4662 Oct 1987 Oct 1987	6.0	1.2*	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N40	B	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	3.0	3.0	3.0	3.0	3.0
4	Revised: Jan 2012 (3rd version)	<b>Kracie Choreito Extract Fine Granules</b> Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)4214 Jul 2007 Jul 2007	6.0	2.5	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-40 3.0g EK-40 2.0g	A	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	3.0	3.0	3.0	3.0	3.0
5	Revised: Apr 2016 (3rd version)	<b>Matsuura Choreito Extract Granules</b> Matsuura Yakugyo Co., Ltd.	(63AM)406 Jul 1988 Jul 1988	6.0	0.7*	Hypromellose, Magnesium Stearate, Dextrin, Lactose Hydrate	Granules	40, Matsuura	A	before or between meals, 2 or 3 times	500g 2g x 84 packets 2g x 300 packets	3.0	3.0	3.0	3.0	3.0
6	Oct 2005 (1st version)	<b>OHSUGI Choreito Extract Granules G</b> Ohsugi Pharmaceutical Co., Ltd.	(62AM)0325 Oct 1987 Oct 1987	6.0	0.4*	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-40	A	before or between meals, 2 or 3 times	500g 588g (2.0g x 294 packets) 168g (2.0g x 84 packets)	3.0	3.0	3.0	3.0	3.0
7	Revised: Jun 2020 (5th version)	<b>SANWA Choreito Extract Fine Granules</b> Sanwa Shoyaku Co., Ltd.	(61AM)3631 Oct 1986 Nov 1986	7.5	3.7	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(S-34)	C	before or between meals, 3 times	500g 2.5g x 300 packets	3.0	3.0	3.0	3.0	3.0
8	Revised: Jun 2011 (4th version)	<b>Taikodo Choreito Extract Granules</b> Taikoseido Pharmaceutical Co., Ltd.	(62AM)0178 Oct 1987 Oct 1987	6.0	4.2	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-40	A	before or between meals, 3 times	500g 2.0g x 84 packets 2.0g x 315 packets	3.0	3.0	3.0	3.0	3.0
9	Revised: Aug 2017 (7th version)	<b>TEIKOKU Choreito Extract Granules</b> Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(62AM)0068 Oct 1987 Oct 1987	7.5	2.49	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 40	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	3.0	3.0	3.0	3.0	3.0
10	Revised: Oct 2005 (4th version)	<b>Toyo Choreito Extract Fine Granules</b> Toyo Yakuko Co., Ltd.	(62AMY)0395 Jul 1988 Oct 1988	6.0	4.0	Corn Starch	Fine granules	TY-080	A	on empty stomach, 3 times	500g 2.0g x 252 packets	3.0	3.0	3.0	3.0	3.0
11	Revised: Mar 2013 (5th version)	<b>TSUMURA Choreito Extract Granules for Ethical Use</b> Tsumura & Co.	(61AM)3282 Oct 1986 Oct 1986	7.5	2.5	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/40	D	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	3.0	3.0	3.0

\*: extract weight without Donkey Glue or Gelatin

Table of Indications

A	The relief of decreased urine volume, difficulty in micturition, and micturition pain or feeling of residual urine
B	The following symptoms of those patients with thirst, painful micturition, dysuria, red urine or blood in urine, edema of lower back or legs: Nephritis, nephrosis, bladder catarrh, urethritis, and dysuria due to renal/bladder calculus
C	Cystitis, especially of acute cystitis, nephritis, dry mouth due to renal calculus or urethritis, increased urinary frequency, and micturition pain
D	The following symptoms of those patients who complain of decreased urine volume, dysuria, and oral dryness: Urethritis, nephritis, renal calculus, blennorrhagic inflammation, micturition pain, hematuria, edema of the lower part of the body from the waist down, feeling of residual urine, and diarrhea