

016 Daijokito

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio			
												Immature Orange	Magnolia Bark	Rhubarb	Anhydrous Sodium Sulfate
1	Revised: Oct 2016 (6th version)	<b>Kotaro Daijokito Extract Fine Granules</b> Kotaro Pharmaceutical Co., Ltd.	(61AM)3966 Oct 1986 Oct 1986	6.0	2.3	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminummetasilicate	Fine granules	N133	A	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	2.0	5.0	2.0	0.9
2	Revised: Mar 2013 (5th version)	<b>TSUMURA Daijokito Extract Granules for Ethical Use</b> Tsumura & Co.	(61AM)1163 Oct 1986 Oct 1986	7.5	3.0	JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/133	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	3.0	5.0	2.0	1.3

**Table of Indications**

A	The following symptoms of those patients who have constipation associated with hardness of and a blocked feeling in the abdomen or those patients who have constipation predisposed to obesity: Chronic constipation, acute constipation, hypertension, neurosis, and food poisoning
---	---