

017 Daikenchuto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio			
												Zanthoxylum Fruit	Ginseng	Processed Ginger	Koi
1	Revised: Dec 2012 (9th version)	Kotaro Daikenchuto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3957 Oct 1986 Oct 1986	27.0	2.1*	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N100	A	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g)	2.0	3.0	5.0	20.0
2	Revised: Jun 2016 (11th version)	TSUMURA Daikenchuto Extract Granules for Ethical Use Tsumura & Co.	(61AM)3299 Oct 1986 Oct 1986	15.0	1.25*	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/100	B	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 84 packets 2.5g x 189 packets	2.0	3.0	5.0	10.0

*: extract weight without Koi

Table of Indications

A	The following symptoms of those patients with relaxation in the intestinal walls and gastrointestinal tract, abdominal cold feeling, vomiting, abdominal distension, increased bowel peristalsis, and severe abdominal pain: Gastroptosis, gastric atony, atonic diarrhea, atonic constipation, chronic peritonitis, and abdominal pain
B	Abdominal cold feeling and pain accompanied by abdominal flatulence