

018 Daiobotampito

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio					
												Rhubarb	Moutan Bark	Peach Kernel	Anhydrous Sodium Sulfate	Sodium Sulfate Hydrate	Benincasa Seed
1	Revised: Oct 2016 (6th version)	Kotaro Daiobotampito Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3826 Oct 1986 Oct 1986	6.0	3.8	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminummetasilicate	Fine granules	N33	B	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	2.0	4.0	4.0	1.8	6.0	
2	Revised: Aug 2017 (7th version)	TEIKOKU Daiobotampito Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3680 Jul 1988 Jul 1988	7.5	2.32	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 33	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	2.0	4.0	4.0	4.0	6.0	
3	Revised: Mar 2013 (5th version)	TSUMURA Daiobotampito Extract Granules for Ethical Use Tsumura & Co.	(61AM)3321 Oct 1986 Oct 1986	7.5	3.5	JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/33	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	2.0	4.0	4.0	1.8	6.0	

Table of Indications

A	The following symptoms of those patients with a comparatively strong constitution and lower abdominal pain who are likely to have constipation: Menstrual irregularity, dysmenorrhea, constipation, and hemorrhoid
B	The following symptoms of those patients with cecal tenderness and fecal impaction, hard feces, red-purple or dark red skin, and tendency to congestion or bleeding: Chronic constipation, arteriosclerosis, various disorders due to menstrual irregularity, climacteric disturbance, eczema, urticaria, acne, skin boils, and bladder catarrh