

021 Daisaikotokyodaio

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio						
												Bupleurum Root	Pinellia Tuber	Ginger	Scutellaria Root	Peony Root	Jujube	Immature Orange
1	Revised: Nov 2007 (4th version)	<b>Kotaro Daisaikotokyodaio Extract Fine Granules</b> Kotaro Pharmaceutical Co., Ltd.	(61AM)3950 Oct 1986 Oct 1986	9.0	5.7	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N319	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	6.0	4.0	1.0	3.0	3.0	3.0	2.0
2	Revised: Feb 2016 (3rd version)	<b>SANWA Daisaikokyodaio Extract Fine Granules</b> Sanwa Shoyaku Co., Ltd.	(61AM)3606 Oct 1986 Nov 1986	9.0	5.6	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-30	A	before or between meals, 3 times	500g 3.0g x 300 packets	6.0	4.0	2.0	3.0	3.0	3.0	2.0
3	Revised: Oct 2018 (4th version)	<b>SANWA Daisaikokyodaio Extract Fine Granules</b> Sanwa Shoyaku Co., Ltd. Ohsugi Pharmaceutical Co., Ltd.	(61AM)3606 Oct 1986 Nov 1986	9.0	5.6	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(SG-148)	A	before or between meals, 3 times	500g 882g (3.0g x 294 packets)	6.0	4.0	2.0	3.0	3.0	3.0	2.0

Table of Indications

A	The following symptoms of those patients who have pressure or pain in the chest or flank, diarrhea, shoulder stiffness and anorexia, etc.: Hepatitis, cholecystitis, cholelithiasis, gastrointestinal catarrh, insomnia, intercostal neuralgia, arteriosclerosis, and hypertension
B	The following symptoms of those patients with epigastric tenderness and pain or pressure in the chest or flank sometimes accompanied by tinnitus, shoulder stiffness, fatigue, and decreased appetite without constipation: Hypertension, arteriosclerosis, gastrointestinal disease, bronchial asthma, jaundice, cholelithiasis, cholecystitis, insomnia, neurasthenia, impotence, pleurisy, hemorrhoids, and hemiplegia