

022 Eppikajutsuto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio					
												Glycyrrhiza	Ginger	Gypsum	Rhizome	Atractylodes Lancea	Jujube
1	Revised: Oct 2010 (5th version)	JPS Eppikajutsuto Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd.	61AM-4759 Oct 1987 Oct 1987	7.5	4.0	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-28)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	2.0	1.0	8.0	4.0	3.0	6.0
2	Revised: Oct 2010 (4th version)	JPS Eppikajutsuto Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd. Ohsugi Pharmaceutical Co., Ltd.	(61AM)4759 Oct 1987 Oct 1987	7.5	4.0	Lactose Hydrate, Magnesium Stearate, Sucrose Esters of Fatty Acids	Granules	(SG-28)	A	before or between meals, 2 or 3 times	500g 735g (2.5g x 294 packets)	2.0	1.0	8.0	4.0	3.0	6.0
3	Revised: Oct 2010 (7th version)	Kotaro Eppikajutsuto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3965 Oct 1986 Oct 1986	9.0	6.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N28	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	2.0	0.8	8.0	4.0	3.0	6.0
4	Revised: Oct 2010 (6th version)	TSUMURA Eppikajutsuto Extract Granules for Ethical Use Tsumura & Co.	(61AM)3304 Oct 1986 Oct 1986	7.5	3.25	JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/28	C	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	2.0	1.0	8.0	4.0	3.0	6.0

Table of Indications

A	The following symptoms of those patients with edema and decreased urinary output: Edema in the early phase of nephritis and nephrosis, edema in beriberi, gonarthrosis, articular rheumatism, acute conjunctivitis, phlyctenular keratoconjunctivitis, pterygium, and eczema
B	The following symptoms of those patients with thirst, severe edema or blisters, decreased urinary output or pollakiuria or excessive secretion: Nephritis, nephrosis, eczema, and beriberi
C	The following symptoms of those patients with edema, diaphoresis, and difficulty in micturition: Nephritis, nephrosis, beriberi, articular rheumatism, nocturnal enuresis, and eczema