

024 Goreisan

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio													
												Albion Tablet	Powdered Albion Tablet	Polyporus Sclerotium	Sclerotium	Poria Sclerotium	Powdered Polyporus Sclerotium	Powdered Poria Sclerotium	Powdered Amanita muscaria Sclerotium	Powdered Amanita muscaria Sclerotium	Amomum Bark	Amomum Twig	Powdered Cinnamon Bark		
1	Revised: Feb 2014 (6th version)	Honzo Goreisan Granules-R Honzo Pharmaceutical Co., Ltd.	(55AM)603 Aug 1981 Aug 1981	5.0	0*	Lactose Hydrate, Carmellose Calcium, Magnesium Aluminometasilicate	Granules	H17	A-1	before meals, 2 times	2.5g x 60 packets	/	0.8	/	0.6	/	0.6	/	0.6	/	0.6	/	/	/	0.4
2	Revised: Feb 2014 (4th version)	JPS Goreisanryo Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd.	61AM-4661 Oct 1987 Oct 1987	7.5	2.4	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-17)	A-1	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	/	6.0	/	4.5	/	4.5	/	4.5	/	4.5	/	3.0	/	/
3	Revised: Feb 2014 (4th version)	JPS Goreisanryo Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd. Ohsumi Pharmaceutical Co., Ltd.	(61AM)4661 Oct 1987 Oct 1987	7.5	2.4	Lactose Hydrate, Magnesium Stearate, Sucrose Esters of Fatty Acids	Granules	(SG-17)	A-1	before or between meals, 2 or 3 times	500g 735g (2.5g x 294 packets)	/	6.0	/	4.5	/	4.5	/	4.5	/	4.5	/	3.0	/	/
4	Revised: Dec 2016 (4th version)	JUNKOU Goreisanryo FC Extract Fine Granules for Ethical Use Kohwayakutuu Inc. Ohsumi Pharmaceutical Co., Ltd.	16100AMY00365000 Oct 1986 Oct 1986	4.5	2.25	Corn Starch, Lactose Hydrate	Fine granules	FC 17	A-2	before or between meals, 2 or 3 times	252g (1.5g x 168 packets)	/	6.0	/	4.5	/	4.5	/	4.5	/	4.5	/	3.0	/	/
5	Revised: Feb 2014 (6th version)	Kotaro Goreisanryo Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)4229 Oct 1986 Oct 1986	6.0	3.2	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N17	B	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	/	6.0	/	4.5	/	4.5	/	4.5	/	4.5	/	2.5	/	/
6	Revised: Feb 2014 (3rd version)	Kracie Goreisanryo Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)4218 Jul 2007 Jul 2007	6.0	2.0	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-17 3.0g EK-17 2.0g	A-1	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	/	5.0	/	3.0	/	3.0	/	3.0	/	3.0	/	2.0	/	/
7	Revised: Feb 2014 (3rd version)	Kracie Goreisanryo Extract Tablets Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00646000 Jul 2007 Jul 2007	5.94 (18 tablets)	2.3	JP Magnesium Stearate, JP Talc, JP Carmellose Calcium, JP Light Anhydrous Silicic Acid, JP Lactose Hydrate, JP Microcrystalline Cellulose, Magnesium Hydroxide-Aluminium Hydroxide Co-Precipitate	Tablets	(EKT-17)	A-1	before or between meals, 2 or 3 times	252 tablets (3 tablets x 6 packets x 14 sheets) 1,764 tablets (3 tablets x 6 packets x 98 sheets)	/	5.0	/	3.0	/	3.0	/	3.0	/	3.0	/	2.0	/	/
8	Revised: Apr 2016 (6th version)	Matsura Goreisanryo Extract Granules Matsura Yakugyo Co., Ltd.	(62AM)585 Oct 1987 Oct 1987	4.5	2.4	Hypromellose, Magnesium Stearate, Lactose Hydrate, Dextrin, Corn Starch	Granules	17, Matsura	A-1	before or between meals, 2 or 3 times	500g 1.5g x 84 packets 1.5g x 300 packets	/	5.0	/	3.0	/	3.0	/	3.0	/	3.0	/	2.0	/	/
9	Revised: Feb 2014 (3rd version)	SANWA Goreisanryo Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3629 Oct 1986 Nov 1986	7.5	3.8	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-33	C	before or between meals, 3 times	500g 2.5g x 300 packets	/	6.0	/	4.5	/	4.5	/	4.5	/	4.5	/	3.0	/	/
10	Revised: Feb 2014 (5th version)	Taikodo Goreisanryo Extract Granules Taikoseido Pharmaceutical Co., Ltd.	(62AM)0177 Oct 1987 Oct 1987	6.0	3.14	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-17	A-1	before or between meals, 3 times	500g 2.0g x 84 packets 2.0g x 315 packets	/	6.0	/	4.5	/	4.5	/	4.5	/	4.5	/	3.0	/	/
11	Revised: Aug 2017 (7th version)	TEIKOKU Goreisan Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3866 Jul 1988 Jul 1988	7.5	1.69	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 17	A-1	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	/	5.0	/	3.0	/	3.0	/	3.0	/	3.0	/	2.0	/	/
12	Revised: Feb 2014 (5th version)	Toyo Goreisanryo Extract Fine Granules Toyo Yakuko Co., Ltd.	(62AM)0397 Jul 1988 Oct 1988	6.0	3.0	Corn Starch	Fine granules	TY-043	A-1	on empty stomach, 3 times	500g 2.0g x 252 packets	/	5.0	/	3.0	/	3.0	/	3.0	/	3.0	/	2.0	/	/
13	Revised: Oct 2014 (6th version)	TSUMURA Goreisan Extract Granules for Ethical Use Tsumura & Co.	(61AM)3287 Oct 1986 Oct 1986	7.5	2.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/17	D	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	/	4.0	/	3.0	/	3.0	/	3.0	/	3.0	/	1.5	/	/

*: It is a crude drug powder (3.0g/day) and not an extract preparation.

Table of Indications

A-1	The following symptoms of those patients with thirst and decreased urine volume who have nausea or vomiting or stomachache or headache or swelling, etc.: Watery diarrhea, acute gastroenteritis (do not use for tenesmus alvi), heat exhaustion, headache, and edema
A-2	The following symptoms of those patients with thirst and decreased urine volume who have nausea or vomiting or stomachache or headache or swelling, etc.: Watery diarrhea, acute gastroenteritis (do not use for tenesmus alvi), heat exhaustion, dull headache, and edema
B	The following symptoms of those patients with thirst and decreased urinary output although drinking water accompanied by headache, heaviness of head, sweating in the head, nausea, vomiting or edema: Acute gastrointestinal catarrh, diarrhea in children and infants, hangover, sun stroke, jaundice, nephritis, nephrosis, and bladder catarrh
C	The following symptoms of those patients with dry mouth, dizziness, headache and edema, etc.: Acute gastrointestinal catarrh, nausea, and nephrosis
D	The following symptoms of those patients with oral dryness and decreased urine volume: Edema, nephrosis, alcoholic hangover, acute gastrointestinal catarrh, diarrhea, nausea, vomiting, dizziness, water retention in the stomach, headache, uremia, heatstroke, and diabetes mellitus