

027 Goshakusan

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio																	
												Portia Sclerotium	Rhizome	Atractylodes Lancea	Atractylodes Rhizome	Citrus Unshiu Peel	Pinellia Tuber	Japanese Angelica Root	Peony Root	Cnidium Rhizome	Magnolia Bark	Angelica Dahurica Root	Orange Fruit	Immature Orange	Platycodon Root	Processed Ginger	Ginger	Cinnamon Bark	Ephedra Herb
1	Revised: May 2008 (6th version)	Kotaro Goshakusan Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3809 Oct 1986 Oct 1986	9.0	5.5	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N63	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	2.0	2.0	2.0	2.0	2.0	2.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	
2	Revised: Aug 2017 (6th version)	TEIKOKU Goshakusan Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3748 Oct 1987 Oct 1987	7.5	3.12	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 63	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	2.0	3.0	2.0	2.0	2.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	0.3	1.0	1.0	1.0	1.0
3	Revised: Nov 2007 (5th version)	TSUMURA Goshakusan Extract Granules for Ethical Use Tsumura & Co.	(61AM)1131 Oct 1986 Oct 1986	7.5	4.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/63	A	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	2.0	3.0	2.0	2.0	2.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0

Table of Indications

A	The following symptoms that take a chronic course without severe symptoms: Gastroenteritis, low back pain, neuralgia, arthralgia, menses painful, headache, oversensitivity to cold, climacteric disturbance, and common cold
B	The following symptoms of those patients with poor circulation, fatigability, and weak stomach: Gastritis, gastric atony, gastroparesis, lower back pain, sciatica, rheumatism, gynecological disorder, and beriberi