

031 Hangebyakujutsutemmato

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Dosage and Administration (Adult)	Packaging	Component Ratio													
												Pinellia Tuber	Arctostaphylos Rhizome	Arctostaphylos Lancea Rhizome	Citrus Unshiu Peel	Portia Selenitum	Malt	Gastrodia Tuber	Ginger	Shinkku	Astragalus Root	Ginseng	Alisma Tuber	Phellodendron Bark	Processed Ginger
1	Revised: Jul 2013 (6th version)	Kotaro Hangebyakujutsutemmato Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3835 Oct 1986 Oct 1986	9.0	6.2	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminummetasilicate	Fine granules	N37	C	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	3.0	3.0	3.0	3.0	3.0	2.0	2.0	0.5	2.0	1.5	1.5	1.5	1.0	1.0
2	Revised: Oct 2013 (3rd version)	Kracie Hangebyakujutsutemmato Extract Fine Granules Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00647000 Jul 2007 Jul 2007	7.5	4.7	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-37 3.75g EK-37 2.5g	B	before or between meals, 2 or 3 times	3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g	3.0	3.0	3.0	3.0	3.0	2.0	2.0	0.65	/	1.5	1.5	1.5	1.0	/
3	Revised: Sep 2017 (4th version)	SANWA Hangebyakujutsutemmato Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3630 Oct 1986 Nov 1986	7.5	4.9	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-37	A	before or between meals, 3 times	500g 2.5g x 300 packets	3.0	3.0	/	3.0	3.0	2.0	2.0	0.5	2.0	1.5	1.5	1.5	1.0	1.0
4	Revised: Sep 2014 (4th version)	SANWA Hangebyakujutsutemmato Extract Fine Granules Sanwa Shoyaku Co., Ltd. JPS Pharmaceutical Co., Ltd.	(61AM)3630 Oct 1986 Nov 1986	7.5	4.9	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(J-37)	A	before or between meals, 3 times	500g 750g (2.5g x 300 packets)	3.0	3.0	/	3.0	3.0	2.0	2.0	0.5	2.0	1.5	1.5	1.5	1.0	1.0
5	Revised: Nov 2018 (5th version)	SANWA Hangebyakujutsutemmato Extract Fine Granules Sanwa Shoyaku Co., Ltd. Ohsugi Pharmaceutical Co., Ltd.	(61AM)3630 Oct 1986 Nov 1986	7.5	4.9	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(SG-37)	A	before or between meals, 3 times	500g 735g (2.5g x 294 packets)	3.0	3.0	/	3.0	3.0	2.0	2.0	0.5	2.0	1.5	1.5	1.5	1.0	1.0
6	Revised: May 2007 (4th version)	TSUMURA Hangebyakujutsutemmato Extract Granules for Ethical Use Tsumura & Co.	(61AM)3258 Oct 1986 Oct 1986	7.5	4.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/37	B	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	/	3.0	3.0	2.0	2.0	0.5	/	1.5	1.5	1.5	1.0	1.0

Table of Indications

A	The following symptoms of those patients with chronic poor digestion, cold lower limbs, occasional headache, and dizziness with vomiting when severe: or patients with languid extremities and lethargy after meals, and frequent substernal splashing sound: Gastric atony, gastropnoia, gastric neurosis, and hypotension
B	Cold lower limbs, dizziness, and headache in patients with a weak gastrointestinal tract
C	The following symptoms of those patients with poor circulation and atonic constitution, fatigability, headache, heaviness of head, dizziness, shoulder stiffness sometimes accompanied by nausea and vomiting: Gastric atony, weak stomach, headache and dizziness due to hypotension