

033 Hangeshashinto

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio						
												Pueraria Tuber	Scutellaria Root	Processed Ginger	Ginger	Ginseng	Glycyrrhiza	Jujube
1	Revised: Jan 2017 (6th version)	Honzo Hangeshashinto Extract Granules-M Honzo Pharmaceutical Co., Ltd.	(61AM)4144 Oct 1986 Oct 1986	7.5	3.4	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminummetasilicate, Magnesium Stearate	Granules	H14	A	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	5.0	2.5	2.5	2.5	2.5	2.5	1.0
2	Revised: Jan 2017 (7th version)	JPS Hangeshashinto Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd.	61AM-4767 Oct 1987 Oct 1987	7.5	4.6	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-14)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	5.0	2.5	2.5	2.5	2.5	2.5	1.0
3	Revised: Jan 2017 (4th version)	JUNKOU Hangeshashinto FC Extract Fine Granules for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00422000 Oct 1986 Oct 1986	6.0	3.2	Corn Starch, Lactose Hydrate	Fine granules	FC 14	A	before or between meals, 2 or 3 times	168g (2.0g x 84 packets)	6.0	3.0	3.0	3.0	3.0	3.0	1.0
4	Revised: Jan 2017 (7th version)	Kotaro Hangeshashinto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3834 Oct 1986 Oct 1986	7.5	5.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminummetasilicate	Fine granules	N14	B	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g) 2.5g x 42 packets (105g)	5.0	2.5	2.5	2.5	2.5	2.5	1.0
5	Revised: Jan 2017 (4th version)	Kracie Hangeshashinto Extract Fine Granules Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00639000 Jul 2007 Jul 2007	6.0	3.8	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-14 3.0g EK-14 2.0g	A	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	5.0	2.5	2.5	2.5	2.5	2.5	1.0
6	Revised: Oct 2013 (3rd version)	Kracie Hangeshashinto Extract Tablets Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00645000 Jul 2007 Jul 2007	5.94 (18 tablets)	3.8	JP Sodium Lauryl Sulfate, JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Lactose Hydrate, JP Microcrystalline Cellulose, JP Croscarmellose Sodium	Tablets	(EKT-14)	A	before or between meals, 2 or 3 times	252 tablets (3 tablets x 6 packets x 14 sheets) 1,764 tablets (3 tablets x 6 packets x 98 sheets)	5.0	2.5	2.5	2.5	2.5	2.5	1.0
7	Revised: Jan 2017 (6th version)	Matsuura Hangeshashinto Extract Granules Matsuura Yakugyo Co., Ltd.	(61AM)4221 Oct 1986 Oct 1986	6.0	viscous extract 7.3*	Hypromellose, Magnesium Stearate, Lactose Hydrate, Dextrin, Corn Starch	Granules	14, Matsuura	A	before or between meals, 2 or 3 times	500g 2g x 84 packets 2g x 300 packets	5.0	2.5	2.5	2.5	2.5	2.5	1.0
8	Revised: Jan 2017 (2nd version)	OHSUGI Hangeshashinto Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3889 Oct 1986 Oct 1986	7.5	3.4	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-14	A	before or between meals, 2 or 3 times	500g 735g (2.5g x 294 packets) 210g (2.5g x 84 packets)	5.0	2.5	2.5	2.5	2.5	2.5	1.0
9	Revised: Mar 2018 (6th version)	SANWA Hangeshashinto Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3598 Oct 1986 Nov 1986	7.5	4.9	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-18	C	before or between meals, 3 times	500g 2.5g x 300 packets	5.0	2.5	2.5	2.5	2.5	2.5	1.0
10	Revised: Jan 2017 (6th version)	Taikodo Hangeshashinto Extract Granules Taikoseido Pharmaceutical Co., Ltd.	(61AM)3911 Oct 1986 Nov 1986	6.0	3.12	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-14	A	before or between meals, 3 times	500g 2.0g x 84 packets 2.0g x 315 packets	5.0	2.5	2.5	2.5	2.5	2.5	1.0
11	Revised: Aug 2017 (9th version)	TEIKOKU Hangeshashinto Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3697 Oct 1987 Oct 1987	9.0	4.31	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 14	A	before meals, 3 times	500g 3.0g x 42 packets 3.0g x 252 packets	5.0	2.5	2.5	2.5	2.5	2.5	1.0
12	Revised: Jan 2017 (6th version)	Toyo Hangeshashinto Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AM)0403 Oct 1986 Dec 1986	6.0	3.6	Corn Starch	Fine granules	TY-094	A	on empty stomach, 3 times	500g 2.0g x 252 packets	5.0	2.5	2.5	2.5	2.5	2.5	1.0
13	Revised: Jan 2017 (9th version)	TSUMURA Hangeshashinto Extract Granules for Ethical Use Tsumura & Co.	(61AM)3257 Oct 1986 Oct 1986	7.5	4.5	JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids	Granules	TSUMURA/14	A	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	5.0	2.5	2.5	2.5	2.5	2.5	1.0

*: correspond to 2.9g of dry extract

Table of Indications

A	The following symptoms of those patients with blocked feeling in the stomach pit and occasional nausea, vomiting, anorexia, borborygmus, and a tendency to loose stools or diarrhea: Acute or chronic gastrointestinal catarrh, fermentative diarrhea, dyspepsia, gastroptosis, nervous gastritis, gastrasthenia, hangover, belching, heartburn, stomatitis, and neurosis
B	The following symptoms of those patients with blocked feeling in the stomach, nausea, and vomiting accompanied by decreased appetite, coated tongue, feeling of residual fluid in the stomach, borborygmus, diarrhea, and soft stools or mucous stools: Acute or chronic gastrointestinal catarrh, fermentative diarrhea, dyspepsia, stomatitis, and morning sickness
C	The following symptoms of those patients with blocked feeling in the stomach, nausea, vomiting, coated tongue, feeling of residual fluid in the stomach, anorexia, borborygmus, and diarrhea or loose stools: Acute or chronic gastrointestinal catarrh, fermentative diarrhea, dyspepsia, gastroptosis, gastric atony, mild illness or prognosis of gastric and duodenal ulcer, and hyperemesis gravidarum