

034 Heisan

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio					
												Atractylodes Lanceae Rhizome	Magnolia Bark	Citrus Unshiu Peel	Jujube	Glycyrrhiza	Ginger
1	Revised: Jun 2009 (4th version)	Honzo Heisanryo Extract Granules-M Honzo Pharmaceutical Co., Ltd.	(61AM)4164 Mar 1989 Mar 1989	7.5	3.0	Magnesium Aluminometasilicate, Microcrystalline Cellulose, Croscarmellose Sodium, Magnesium Stearate, Light Anhydrous Silicic Acid, Potato Starch, Lactose Hydrate	Granules	H79	A	before or between meals, 2 or 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	4.0	3.0	3.0	2.0	1.0	0.5
2	Revised: Feb 2008 (4th version)	Kotaro Heisan Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3958 Oct 1986 Oct 1986	6.0	4.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N79	B	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	4.0	3.0	3.0	2.0	1.0	0.5
3	Oct 2005 (1st version)	OHSUGI Heisanryo Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3897 Oct 1986 Oct 1986	7.5	2.9	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-79	A	before or between meals, 2 or 3 times	500g 735g (2.5g x 294 packets) 210g (2.5g x 84 packets)	4.0	3.0	3.0	2.0	1.0	0.5
4	Revised: Aug 2017 (6th version)	TEIKOKU Heisan Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3687 Jul 1988 Jul 1988	7.5	1.9	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 79	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	4.0	3.0	3.0	2.0	1.0	0.5
5	Revised: Mar 2013 (5th version)	TSUMURA Heisan Extract Granules for Ethical Use Tsumura & Co.	(61AM)3273 Oct 1986 Oct 1986	7.5	3.25	JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/79	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	4.0	3.0	3.0	2.0	1.0	0.5

Table of Indications

A	The following symptoms of those patients with a heavy stomach feeling and a tendency to dyspepsia: Acute or chronic esogastritis, gastric atony, dyspepsia, and anorexia
B	The following symptoms of those patients with stomachache accompanied by dyspepsia, abdominal pain, decreased appetite, borborygmus after meals, and tendency to diarrhea: Stomatitis, gastritis, gastric atony, and gastric dilatation