

035 Hochuekkito

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio													
												Ginseng	Atractylodes Rhizome	Atractylodes Lancea Rhizome	Atractylodes Root	Japanese Angelica Root	Citrus Unshiu Peel	Jungho	Bupleurum Root	Glycyrrhiza	Ginger	Fresh Ginger	Processed Ginger	Cinnabark Rhizome	
1	Revised: Jun 2007 (5th version)	Honzo Hochuekkito Extract Granules-M Honzo Pharmaceutical Co., Ltd.	(61AM)4143 Oct 1986 Oct 1986	7.5	5.0	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate	Granules	H41	A	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	4.0	4.0	/	4.0	3.0	2.0	2.0	1.0	1.5	/	/	0.5	0.5	
2	Revised: Jun 2007 (5th version)	JPS Hochuekkito Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd.	62AM-0480 Oct 1987 Oct 1987	7.5	4.7	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-41)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	4.0	/	4.0	4.0	3.0	2.0	2.0	1.0	1.5	0.5	/	/	0.5	0.5
3	Revised: Nov 2016 (4th version)	JUNKOU Hochuekkito FC Extract Fine Granules for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00380000 Oct 1986 Oct 1986	7.5	4.9	Corn Starch, Lactose Hydrate	Fine granules	FC 41	A	before or between meals, 2 or 3 times	210g (2.5g x 84 packets)	4.0	4.0	/	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
4	Revised: Oct 2016 (2nd version)	JUNKOU Hochuekkito FC Tablet for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	16200AMY00036000 Dec 2014 Mar 2015	7.5 (18 tablets)	4.9	Corn Starch, Lactose Hydrate, Talc, Magnesium Stearate	Tablets	FC 41T	A	before or between meals, 2 or 3 times	504Tablets(3Tablets x168packets)	4.0	4.0	/	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
5	Revised: Nov 2007 (6th version)	Kotaro Hochuekkito Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3949 Oct 1986 Oct 1986	12.0	7.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N41	B	before or between meals, 2 or 3 times	500g 4.0g x 147 packets (588g) 4.0g x 42 packets (168g)	4.0	4.0	/	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
6	Revised: Jul 2010 (2nd version)	Kracie Hochuekkito Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3511 Jul 2007 Jul 2007	7.5	6.4	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Light Anhydrous Silicic Acid, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-41 3.75g EK-41 2.5g	A	before or between meals, 2 or 3 times	3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g	4.0	4.0	/	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
7	Oct 2005 (1st version)	OHSUGI Hochuekkito Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3898 Oct 1986 Oct 1986	12.0	6.2	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-41	A	before or between meals, 2 or 3 times	500g 1,176g (4.0g x 294 packets) 336g (4.0g x 84 packets)	4.0	4.0	/	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
8	Revised: Aug 2015 (5th version)	SANWA Hochuekkito Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3616 Oct 1986 Nov 1986	9.0	5.3	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-12	C	before or between meals, 3 times	500g 3g x 300 packets	4.0	4.0	/	3.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
9	Revised: Oct 2010 (6th version)	Taikodo Hochuekkito Extract Granules Taikoseido Pharmaceutical Co., Ltd.	(61AM)3913 Oct 1986 Nov 1986	7.5	5.55	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-41	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	4.0	4.0	/	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
10	Revised: Mar 2011 (5th version)	Taikodo Hochuekkito Extract Powder Taikoseido Pharmaceutical Co., Ltd.	(61AM)3860 Oct 1986 Nov 1986	7.5	5.55	JP Lactose Hydrate, JP Magnesium Stearate	Powders	TM-41P	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	4.0	4.0	/	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0
11	Revised: Jan 2007 (6th version)	TEIKOKU Hochuekkito Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Ohki Pharmaceutical Co., Ltd.	(61AM)3701 Oct 1987 Oct 1987	7.5	3.57	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	41/OHKI	A	before meals, 3 times	2.5g x 210 packets	4.0	4.0	/	4.0	3.0	2.0	2.0	1.0	1.5	0.5	/	/	0.5	0.5
12	Revised: Aug 2017 (8th version)	TEIKOKU Hochuekkito Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3701 Oct 1987 Oct 1987	7.5	3.57	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 41	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	4.0	4.0	/	4.0	3.0	2.0	2.0	1.0	1.5	0.5	/	/	0.5	0.5
13	Revised: Jul 2005 (6th version)	Toyo Hochuekkito Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AMY)0477 Oct 1987 Jan 1988	7.5	4.8	Corn Starch	Fine granules	TY-101	A	on empty stomach, 3 times	500g 2.5g x 252 packets	4.0	4.0	/	3.0	3.0	2.0	2.0	2.0	1.5	/	2.0	/	1.0	1.0
14	Revised: Oct 2014 (11th version)	TSUMURA Hochuekkito Extract Granules for Ethical Use Tsumura & Co.	(61AM)1164 Oct 1986 Oct 1986	7.5	5.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/41	D	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	4.0	/	4.0	4.0	3.0	2.0	2.0	2.0	1.5	0.5	/	/	1.0	1.0

Table of Indications

A	The following symptoms of those patients who are dispirited, having poor gastrointestinal function and are likely to get tired: Infirm constitution, fatigue and malaise, declined constitution after disease, anorexia, and night sweats
B	The following symptoms of those patients with decreased gastrointestinal function, fatigue, and malaise, or headache, chills, night sweating, and atonic hemorrhage: Enhancement of physical strength in patients with tuberculous disease or after disease, weak stomach, decreased weight in summer, delicate constitution, hypotension, scrofulous constitution, hemorrhoids, and anal prolapse
C	The following symptoms of those patients who have a delicate constitution, anemia, reduced gastrointestinal functions, fatigue and malaise, anorexia, or perspiration during sleep, etc.: Debility after disease or surgery, reinforcement of physical strength after chest disease, anemia, hypotension, summer emaciation, dyspepsia, weak gastrointestinal functions, and hyperhidrosis
D	The following symptoms/conditions of patients having delicate constitution, reduced digestive functions, and severe fatigability of limbs: Summer emaciation, reinforcement of physical strength after illness, tuberculosis, anorexia, gastroptosis, cold, hemorrhoid, anal prolapse, uterine prolapse, impotence, hemiplegia, and hyperhidrosis