

035 Hochuekkito

| No. | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | | | | | | | | | |
|-----|--|--|---|-------------------------|--|---|----------------|-----------------------------------|-------------|---|--|-----------------|------------------------|----------------------------------|--------------------|---------------------------|-----------------------|--------|-------------------|-------------|--------|-----------------|---------------------|--------------------|-----|
| | | | | | | | | | | | | Ginseng | Anacycloles Rhizome | Anacycloles Lancea Rhizome | Astringent Root | Japanese Angelica Root | Citrus Unshiu Peel | Jujube | Bupleurum Root | Glycyrrhiza | Ginger | Fresh Ginger | Processed Ginger | Cnidium Rhizome | |
| 1 | Revised: Jun 2007 (5th version) | Honzo Hochuekkito Extract Granules-M Honzo Pharmaceutical Co., Ltd. | (61AM)4143 Oct 1986 Oct 1986 | 7.5 | 5.0 | Lactose Hydrate, Microcrystalline Cellulose, Magnesium Akuminometasilicate, Magnesium Stearate | Granules | H41 | A | before or between meals, 3 times | 500g 2.5g x 294 packets 2.5g x 42 packets | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 1.0 | 1.5 | / | / | 0.5 | 0.5 | |
| 2 | Revised: Jun 2007 (5th version) | JPS Hochuekkito Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd. | 62AM-0480 Oct 1987 Oct 1987 | 7.5 | 4.7 | Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate | Granules | (J-41) | A | before or between meals, 2 or 3 times | 500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets) | 4.0 | / | 4.0 | 4.0 | 3.0 | 2.0 | 2.0 | 1.0 | 1.5 | 0.5 | / | / | 0.5 | 0.5 |
| 3 | Revised: Nov 2016 (4th version) | JUNKOU Hochuekkito FC Extract Fine Granules for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd. | 16100AMY00380000 Oct 1986 Oct 1986 | 7.5 | 4.9 | Corn Starch, Lactose Hydrate | Fine granules | FC 41 | A | before or between meals, 2 or 3 times | 210g (2.5g x 84 packets) | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 4 | Revised: Oct 2016 (2nd version) | JUNKOU Hochuekkito FC Tablet for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd. | 16200AMY0036000 Dec 2014 Mar 2015 | 7.5 (18 tablets) | 4.9 | Corn Starch, Lactose Hydrate, Tak, Magnesium Stearate | Tablets | FC 41T | A | before or between meals, 2 or 3 times | 504Tablets(3Tablets x168packets) | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 5 | Revised: Nov 2007 (6th version) | Kotaro Hochuekkito Extract Fine Granules Kotaro Pharmaceutical Co., Ltd. | (61AM)3949 Oct 1986 Oct 1986 | 12.0 | 7.0 | Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate | Fine granules | N41 | B | before or between meals, 2 or 3 times | 500g 4.0g x 147 packets (588g) 4.0g x 42 packets (168g) | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 6 | Revised: Jul 2010 (2nd version) | Kracie Hochuekkito Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd. | (61AM)3511 Jul 2007 Jul 2007 | 7.5 | 6.4 | JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Light Anhydrous Silicic Acid, JP Lactose Hydrate, Hydrated Silicon Dioxide | Fine granules | KB-41 3.75g EK-41 2.5g | A | before or between meals, 2 or 3 times | 3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 7 | Oct 2005 (1st version) | OHSUGI Hochuekkito Extract Granules G Ohsugi Pharmaceutical Co., Ltd. | (61AM)3898 Oct 1986 Oct 1986 | 12.0 | 6.2 | Lactose Hydrate, Corn Starch, Magnesium Stearate | Granules | SG-41 | A | before or between meals, 2 or 3 times | 500g 1,176g (4.0g x 294 packets) 336g (4.0g x 84 packets) | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 8 | Revised: Aug 2015 (5th version) | SANWA Hochuekkito Extract Fine Granules Sanwa Shoyaku Co., Ltd. | (61AM)3616 Oct 1986 Nov 1986 | 9.0 | 5.3 | Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid | Fine granules | S-12 | C | before or between meals, 3 times | 500g 3g x 300 packets | 4.0 | 4.0 | / | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 9 | Revised: Oct 2010 (6th version) | Taikodo Hochuekkito Extract Granules Taikoseido Pharmaceutical Co., Ltd. | (61AM)3913 Oct 1986 Nov 1986 | 7.5 | 5.55 | JP Lactose Hydrate, JP Magnesium Stearate | Granules | TM-41 | A | before or between meals, 3 times | 500g 2.5g x 84 packets 2.5g x 252 packets | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 10 | Revised: Mar 2011 (5th version) | Taikodo Hochuekkito Extract Powder Taikoseido Pharmaceutical Co., Ltd. | (61AM)3860 Oct 1986 Nov 1986 | 7.5 | 5.55 | JP Lactose Hydrate, JP Magnesium Stearate | Powders | TM-41P | A | before or between meals, 3 times | 500g 2.5g x 84 packets 2.5g x 252 packets | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |
| 11 | Revised: Aug 2017 (8th version) | TEIKOKU Hochuekkito Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd. | (61AM)3701 Oct 1987 Oct 1987 | 7.5 | 3.57 | Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate | Granules | TEIKOKU 41 | A | before meals, 3 times | 500g 2.5g x 42 packets 2.5g x 252 packets | 4.0 | 4.0 | / | 4.0 | 3.0 | 2.0 | 2.0 | 1.0 | 1.5 | 0.5 | / | / | 0.5 | 0.5 |
| 12 | Revised: Jul 2005 (6th version) | Toyo Hochuekkito Extract Fine Granules Toyo Yakuko Co., Ltd. | (61AMY)0477 Oct 1987 Jan 1988 | 7.5 | 4.8 | Corn Starch | Fine granules | TY-101 | A | on empty stomach, 3 times | 500g 2.5g x 252 packets | 4.0 | 4.0 | / | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | / | 2.0 | / | 1.0 | 1.0 |
| 13 | Revised: Oct 2014 (11th version) | TSUMURA Hochuekkito Extract Granules for Ethical Use Tsumura & Co. | (61AM)1164 Oct 1986 Oct 1986 | 7.5 | 5.0 | JP Magnesium Stearate, JP Lactose Hydrate | Granules | TSUMURA/41 | D | before or between meals, 2 or 3 times | 500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets | 4.0 | / | 4.0 | 4.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.5 | 0.5 | / | / | 1.0 | 1.0 |

Table of Indications

| | |
|---|---|
| A | The following symptoms of those patients who are dispirited, having poor gastrointestinal function and are likely to get tired: Infirm constitution, fatigue and malaise, declined constitution after disease, anorexia, and night sweats |
| B | The following symptoms of those patients with decreased gastrointestinal function, fatigue, and malaise, or headache, chills, night sweating, and atonic hemorrhage: Enhancement of physical strength in patients with tuberculous disease or after disease, weak stomach, decreased weight in summer, delicate constitution, hypotension, scrofulous constitution, hemorrhoids, and anal prolapse |
| C | The following symptoms of those patients who have a delicate constitution, anemia, reduced gastrointestinal functions, fatigue and malaise, anorexia, or perspiration during sleep, etc.: Debility after disease or surgery, reinforcement of physical strength after chest disease, anemia, hypotension, summer emaciation, dyspepsia, weak gastrointestinal functions, and hyperhidrosis |
| D | The following symptoms/conditions of patients having delicate constitution, reduced digestive functions, and severe fatigability of limbs: Summer emaciation, reinforcement of physical strength after illness, tuberculosis, anorexia, gastroptosis, cold, hemorrhoid, anal prolapse, uterine prolapse, impotence, hemiplegia, and hyperhidrosis |