

038 Ireito

| | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | | | | | |
|---|--|---|---|----------------------------|--|---|----------------|-----------------------------------|-------------|---|---|----------------------|---------------------|----------------------|------------------|--------------------------|-------------------------|-----------------|---------------------|------------------|--------|
| | | | | | | | | | | | | Arctiodes Rhizome | Arctiodes Lancea | Arctiodes Rhizome | Magnolia Bark | Citrus Unshiu Peel | Polyporus Sclerotium | Alisma Tuber | Poria Sclerotium | Cinnamon Bark | Jujube |
| 1 | Revised: Mar 2013 (5th version) | TSUMURA Ireito Extract Granules for Ethical Use Tsumura & Co. | (61AM)1148 Oct 1986 Oct 1986 | 7.5 | 4.25 | JP Magnesium Stearate, JP Lactose Hydrate | Granules | TSUMURA/115 | A | before or between meals, 2 or 3 times | 500g 2.5g x 42 packets 2.5g x 189 packets | 2.5 | 2.5 | 2.5 | 2.5 | 2.5 | 2.5 | 2.0 | 1.5 | 1.5 | 1.0 |

Table of Indications

| | |
|---|--|
| A | The following symptoms of those patients with watery diarrhea and vomiting accompanied by thirst and decreased urine volume: Food poisoning, heat stroke, cold abdomen, acute gastroenteritis, and abdominal pain |
|---|--|