

046 Juzentaihoto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio									
												Ginseng	Astragalus Root	Atractylodes Rhizome	Atractylodes Lancoa Rhizome	Poria Sclerotium	Japanese Angelica Root	Peony Root	Rehmannia Root	Cnidium Rhizome	Cinnamon Bark
1	Revised: Jul 2011 (4th version)	Honzo Juzentaihoto Extract Granules-M Honzo Pharmaceutical Co., Ltd.	(61AM)4162 Oct 1986 Oct 1986	9.0	6.1	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate	Granules	H48	A	before or between meals, 3 times	500g 3.0g x 294 packets 3.0g x 42 packets	2.5	2.5	3.5	3.5	3.5	3.0	3.5	3.0	3.0	1.0
2	Revised: May 2017 (4th version)	JUNKOU Juzentaihoto FC Extract Fine Granules for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00375000 Oct 1986 Oct 1986	7.5	5.35	Corn Starch, Lactose Hydrate	Fine granules	FC 48	A	before or between meals, 2 or 3 times	210g (2.5g x 84 packets)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.0
3	Revised: Jan 2012 (5th version)	Kotaro Juzentaihoto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3811 Oct 1986 Oct 1986	15.0	8.5	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N48	C	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g)	2.5	2.5	3.5	3.5	3.5	3.0	3.5	3.0	3.0	1.0
4	Revised: Jan 2012 (3rd version)	Kracie Juzentaihoto Extract Fine Granules Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00623000 Jul 2007 Jul 2007	7.5	6.2	JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-48 3.75g EK-48 2.5g	A	before or between meals, 2 or 3 times	3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.5
5	Oct 2005 (1st version)	OHSUGI Juzentaihoto Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3884 Oct 1986 Oct 1986	12.0	6.1	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-48	A	before or between meals, 2 or 3 times	500g 1,176g (4.0g x 294 packets) 336g (4.0g x 84 packets)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.5
6	Revised: Jun 2012 (4th version)	SANWA Juzentaihoto Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3605 Oct 1986 Nov 1986	9.0	6.1	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-32	B	before or between meals, 3 times	500g 3g x 300 packets	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.5
7	Revised: Mar 2018 (5th version)	SANWA Juzentaihoto Extract Fine Granules Sanwa Shoyaku Co., Ltd. JPS Pharmaceutical Co., Ltd.	61AM-3605 Oct 1986 Nov 1986	9.0	6.1	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(J-48)	B	before or between meals, 3 times	500g 720g (3.0g x 240 packets)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.5
8	Revised: Aug 2017 (7th version)	TEIKOKU Juzentaihoto Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co.,Ltd.	(61AM)3693 Oct 1987 Oct 1987	9.0	4.36	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 48	A	before meals, 3 times	500g 3.0g x 42 packets 3.0g x 252 packets	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.5
9	Revised: Jul 2005 (5th version)	Toyo Juzentaihoto Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AMY)0392 Oct 1986 Jan 1987	9.0	5.7	Corn Starch	Fine granules	TY-060	A	on empty stomach, 3 times	500g 3.0g x 252 packets	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.5
10	Revised: Oct 2014 (7th version)	TSUMURA Juzentaihoto Extract Granules for Ethical Use Tsumura & Co.	(61AM)1124 Oct 1986 Oct 1986	7.5	5.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/48	A	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	1.5

Table of Indications

A	Declined constitution after recovery from disease, fatigue and malaise, anorexia, perspiration during sleep, cold limbs, and anemia
B	The following symptoms of those patients who have anemia, a pale skin and mucous membrane, malnutrition, emaciation with no appetite and debility: Anemia due to debility (after childbirth, surgery or severe illness) , hypotension, leukemia, anal fistula, caries, debility due to wasting disease, hemorrhage, and anal prolapse
C	The following symptoms of those patients with pale skin and mucosa, muddy complexion, thin body, anemia, anorexia or excessive debility as well as the following symptoms in debility due to wasting disease or after surgery, after childbirth, and general debility: Hypotension, anemia, neurasthenia, fatigue, malaise, weak stomach, and gastroptosis