

047 Kakkonkajutsubuto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio									
												Pueraria Root	Ephedra Herb	Cinnamon Bark	Glycyrrhiza	Peony Root	Jujube	Ginger	Atractylodes Rhizome	Lancea	Processed Aconite Root
1	Revised: Mar 2022 (1st version of new form)	SANWA Kakkonkajutsubuto Extract Fine Granules Sanwa Shoyaku Co., Ltd.	16100AMZ03625000 - Nov 1986	7.5	4.8	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-07	A	before or between meals, 3 times	500g [bottle] 2.5g x 84 packets 2.5g x 300 packets	4.0	3.0	2.0	2.0	2.0	3.0	1.0	3.0	0.5	
2	Revised: Oct 2018 (5th version)	SANWA Kakkonkajutsubuto Extract Fine Granules Sanwa Shoyaku Co., Ltd. Ohsugi Pharmaceutical Co., Ltd.	(61AM)3625 Oct 1986 Nov 1986	7.5	4.8	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(SG-141)	A	before or between meals, 3 times	500g 735g (2.5g x 294 packets)	4.0	3.0	2.0	2.0	2.0	3.0	1.0	3.0	0.5	

Table of Indications

A	The following symptoms of those patients who have rigor, fever, headache, and tension in the nuchal, shoulder and back: Shoulder stiffness, neuralgia in the scapular region, and articular rheumatism in the upper body
---	--