

057 Keishakuchimoto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio									
												Cinnamon Bark	Anemarrhena Rhizome	Glehnia Root and Rhizome	Ginger	Peony Root	Ephedra Herb	Atacyliodes Rhizome	Glycyrrhiza	Processed Aconite Root	
1	Revised: Mar 2022 (1st version of new form)	SANWA Keishakuchimoto Extract Fine Granules Sanwa Shoyaku Co., Ltd.	16100AMZ03607000 - Nov 1986	9.0	5.1	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-10	A	before or between meals, 3 times	500g [bottle] 3g x 300 packets	3.0	3.0	3.0	1.0	3.0	3.0	4.0	1.5	1.0	
2	Revised: Jul 2010 (5th version)	SANWA Keishakuchimoto Extract Fine Granules Sanwa Shoyaku Co., Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3607 Oct 1986 Nov 1986	9.0	5.1	JP Lactose Hydrate, JP Corn Starch, JP Microcrystalline Cellulose, JP Light Anhydrous Silicic Acid, Partly Pregelatinized Starch	Fine granules	EK-180	A	before or between meals, 3 times	500g 3.0g x 42 packets 3.0g x 294 packets	3.0	3.0	3.0	1.0	3.0	3.0	4.0	1.5	1.0	

Table of Indications

A	The following symptoms of those patients with painful joints, emaciation, swelling legs, dizziness, nausea: Neuralgia and articular rheumatism
---	--