

066 Keishikashakuyakudaioto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio					
												Cinnamon Bark	Peony Root	Jujube	Ginger	Glycyrrhiza	Rhubarb
1	Revised: Mar 2013 (5th version)	TSUMURA Keishikashakuyakudaioto Extract Granules for Ethical Use Tsumura & Co.	(61AM)1170 Oct 1986 Oct 1986	7.5	4.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/134	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	4.0	6.0	4.0	1.0	2.0	2.0

Table of Indications

A	The following symptoms of those patients with comparatively deprived constitution having abdominal distension accompanied by stagnation of intestinal contents or abdominal pain: 1. Acute enterocolitis and large intestinal catarrh 2. Habitual constipation, fecal impaction, and tenesmus alvi
---	--