

066 Keishikashakuyakudaioto

|   | Date of Making or Revision of Package Insert | Product Name<br>Manufacturer<br>Distributor  | Approval No.<br>Date of Listing in the NHI Reimbursement Price<br>Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients                      | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult)  | Packaging                                       | Component Ratio |            |        |        |             |         |
|---|--|--|--|----------------------|-------------------------------------|---|-------------|-----------------------------|-------------|--|---|-----------------|------------|--------|--------|-------------|---------|
|   |  |  |  |                      |                                     |   |             |                             |             |  |   | Cinnamon Bark   | Peony Root | Jujube | Ginger | Glycyrrhiza | Rhubarb |
| 1 | Revised: Mar 2013 (5th version)              | <b>TSUMURA Keishikashakuyakudaioto Extract Granules for Ethical Use</b><br>Tsumura & Co. | (61AM)1170<br>Oct 1986<br>Oct 1986   | 7.5                  | 4.0                                 | JP Magnesium Stearate, JP Lactose Hydrate | Granules    | TSUMURA/134                 | A           | before or between meals,<br>2 or 3 times | 500g<br>2.5g x 42 packets<br>2.5g x 189 packets | 4.0             | 6.0        | 4.0    | 1.0    | 2.0         | 2.0     |

**Table of Indications**

|   |  |
|---|--|
| A | The following symptoms of those patients with comparatively deprived constitution having abdominal distension accompanied by stagnation of intestinal contents or abdominal pain:<br>1. Acute enterocolitis and large intestinal catarrh<br>2. Habitual constipation, fecal impaction, and tenesmus alvi |
|---|--|