

068 Keishininjinto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio					
												Cinnamon Bark	Glycyrrhiza	Ginseng	Processed Ginger	Atractylodes Rhizome	Atractylodes Lancea
1	Revised: Jul 2010 (2nd version)	Kracie Keishininjinto Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3571 Jul 2007 Jul 2007	6.0	2.7	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-82 3.0g EK-82 2.0g	A	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	4.0	3.0	3.0	2.0	3.0	
2	Revised: Mar 2013 (5th version)	TSUMURA Keishininjinto Extract Granules for Ethical Use Tsumura & Co.	(61AM)1162 Oct 1986 Oct 1986	7.5	2.5	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/82	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	4.0	3.0	3.0	2.0	3.0	

Table of Indications

A	The following symptoms of those patients with a weak digestive system: Headache, palpitation, chronic gastroenteritis, and gastric atony
---	---