

076 Kyukikyogaito

| | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | | |
|---|--|--|--|----------------------|-------------------------------------|---|---------------|-----------------------------|-------------|--|---|-----------------|-------------|----------------|------------------------|------------|----------------|-------------|
| | | | | | | | | | | | | Cnidium Rhizome | Glycyrrhiza | Artemisia Leaf | Japanese Angelica Root | Peony Root | Rehmannia Root | Donkey Glue |
| 1 | Revised: Sep 2017 (3rd version) | JUNKOU Kyukikyogaito FC Extract Fine Granules for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd. | 16100AMY00416000 Oct 1986 Oct 1986 | 9.0 | 5.6 | Corn Starch, Lactose Hydrate | Fine granules | FC 77 | A | before or between meals, 2 or 3 times | 126g (1.5g x 84 packets) | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 5.0 | 3.0 |
| 2 | Revised: Jun 2013 (5th version) | Kotaro Kyukikyogaito Extract Fine Granules Kotaro Pharmaceutical Co., Ltd. | (61AM)3823 Oct 1986 Oct 1986 | 15.0 | 7.0* | Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate | Fine granules | N77 | B | before or between meals, 2 or 3 times | 500g 2.5g x 189 packets (472.5g) | 3.0 | 3.0 | 3.0 | 4.5 | 4.5 | 6.0 | 3.0 |
| 3 | Revised: Mar 2013 (5th version) | TSUMURA Kyukikyogaito Extract Granules for Ethical Use Tsumura & Co. | (61AM)3300 Oct 1986 Oct 1986 | 9.0 | 6.0 | JP Magnesium Stearate, JP Lactose Hydrate | Granules | TSUMURA/77 | A | before or between meals, 2 or 3 times | 500g 3.0g x 42 packets 3.0g x 189 packets | 3.0 | 3.0 | 3.0 | 4.0 | 4.0 | 5.0 | 3.0 |

*: extract weight without Gelatin

Table of Indications

| | |
|---|--|
| A | Hemorrhoidal bleeding |
| B | The following symptoms of those patients with poor circulation and anemia due to excessive bleeding: Hemorrhoidal hemorrhage, bruise after trauma, hemorrhage after childbirth, and anemia |