

077 Makyokansekitō

|   | Date of Making or Revision of Package Insert | Product Name<br>Manufacturer<br>Distributor   | Approval No.<br>Date of Listing in the NHI Reimbursement Price<br>Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients   | Dosage Form   | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult)  | Packaging  | Component Ratio |                |             |        |
|---|--|---|--|----------------------|-------------------------------------|--|---------------|-----------------------------|-------------|--|--|-----------------|----------------|-------------|--------|
|   |  |   |  |                      |                                     |  |               |                             |             |  |  | Ephedra Herb    | Apricot Kernel | Glycyrrhiza | Gypsum |
| 1 | Revised: Jun 2009 (4th version)              | <b>Honzo Makyokansekitō Extract Granules-M</b><br>Honzo Pharmaceutical Co., Ltd.  | (61AM)4166<br>Apr 1989<br>Mar 1989   | 7.5                  | 1.7                                 | Magnesium Aluminummetasilicate, Microcrystalline Cellulose, Croscarmellose Sodium Type A, Light Anhydrous Silicic Acid, Magnesium Stearate, Potato Starch, Lactose Hydrate | Granules      | H55                         | A           | before or between meals,<br>2 or 3 times | 500g<br>2.5g x 294 packets<br>2.5g x 42 packets                    | 4.0             | 4.0            | 2.0         | 10.0   |
| 2 | Revised: Oct 2016 (3rd version)              | <b>JUNKOU Makyokansekitō FC Extract Fine Granules for Ethical Use</b><br>Kohwayakutuu Inc.<br>Ohsugi Pharmaceutical Co., Ltd. | 16100AMY00382000<br>Oct 1986<br>Oct 1986   | 4.5                  | 2.25                                | Corn Starch, Lactose Hydrate   | Fine granules | FC 55                       | A           | before or between meals,<br>2 or 3 times | 126g (1.5g x 84 packets)   | 4.0             | 4.0            | 2.0         | 10.0   |
| 3 | Revised: May 2008 (6th version)              | <b>Kotaro Makyokansekitō Extract Fine Granules</b><br>Kotaro Pharmaceutical Co., Ltd.   | (61AM)3962<br>Oct 1986<br>Oct 1986   | 6.0                  | 2.2                                 | Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminummetasilicate   | Fine granules | N55                         | B           | before or between meals,<br>2 or 3 times | 500g<br>2.0g x 231 packets (462g)<br>2.0g x 42 packets (84g)       | 4.0             | 4.0            | 2.0         | 10.0   |
| 4 | Revised: Apr 2016 (3rd version)              | <b>Matsuura Makyokansekitō Extract Granules</b><br>Matsuura Yakugyo Co., Ltd.   | (61AM)4043<br>Oct 1987<br>Oct 1987   | 4.5                  | 1.9                                 | Hypromellose, Magnesium Stearate, Lactose Hydrate, Dextrin, Corn Starch  | Granules      | 55, Matsuura                | A           | before or between meals,<br>2 or 3 times | 500g<br>1.5g x 84 packets<br>1.5g x 300 packets                    | 4.0             | 4.0            | 2.0         | 10.0   |
| 5 | Oct 2005 (1st version)                       | <b>OHSUGI Makyokansekitō Extract Granules G</b><br>Ohsugi Pharmaceutical Co., Ltd.  | (61AM)3891<br>Oct 1986<br>Oct 1986   | 4.5                  | 1.5                                 | Lactose Hydrate, Corn Starch, Magnesium Stearate   | Granules      | SG-55                       | A           | before or between meals,<br>2 or 3 times | 500g<br>441g (1.5g x 294 packets)<br>126g (1.5g x 84 packets)      | 4.0             | 4.0            | 2.0         | 10.0   |
| 6 | Revised: Aug 2017 (6th version)              | <b>TEIKOKU Makyokansekitō Extract Granules</b><br>Teikoku Kampo Seiyaku Co., Ltd.<br>Teikoku Seiyaku Co., Ltd.                | (61AM)3682<br>Jul 1988<br>Jul 1988   | 7.5                  | 1.22                                | Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate  | Granules      | TEIKOKU 55                  | A           | before meals,<br>3 times                 | 500g<br>2.5g x 42 packets<br>2.5g x 252 packets                    | 4.0             | 4.0            | 2.0         | 10.0   |
| 7 | Revised: Sep 2007 (5th version)              | <b>TSUMURA Makyokansekitō Extract Granules for Ethical Use</b><br>Tsumura & Co.   | (61AM)3302<br>Oct 1986<br>Oct 1986   | 7.5                  | 1.75                                | JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate   | Granules      | TSUMURA/55                  | A           | before or between meals,<br>2 or 3 times | 500g<br>5kg (500g x 10)<br>2.5g x 42 packets<br>2.5g x 189 packets | 4.0             | 4.0            | 2.0         | 10.0   |

Table of Indications

|   |   |
|---|---|
| A | Infantile asthma and bronchial asthma   |
| B | The following symptoms of those patients with severe cough, sweating in the head at the time of attack accompanied by wheezing and thirst:<br>Bronchitis and bronchial asthma |