

085 Ninjinto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio				
												Ginseng	Glycyrrhiza	Atractylodes Rhizome	Atractylodes Lancea	Processed Ginger
1	Revised: Jun 2009 (4th version)	<b>Honzo Ninjinto Extract Fine Granules</b> Honzo Pharmaceutical Co., Ltd.	(61AM)3744 Oct 1987 Oct 1987	6.0	2.6	Lactose Hydrate, Potato Starch, Magnesium Aluminometasilicate, Hydroxypropyl Starch, Carmellose Calcium, Low Substituted Hydroxypropylcellulose	Fine granules	HG32	A	before or between meals, 2 or 3 times	500g 2.0g x 294 packets 2.0g x 42 packets	3.0	3.0	3.0	3.0	3.0
2	Revised: Nov 2007 (6th version)	<b>Kotaro Ninjinto Extract Fine Granules</b> Kotaro Pharmaceutical Co., Ltd.	(61AM)3813 Oct 1986 Oct 1986	6.0	3.2	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N32	B	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	3.0	3.0	3.0	3.0	3.0
3	Revised: Jul 2010 (2nd version)	<b>Kracie Ninjinto Extract Fine Granules</b> Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00640000 Jul 2007 Jul 2007	6.0	3.0	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-32 3.0g EK-32 2.0g	A	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	3.0	3.0	3.0	3.0	3.0
4	Revised: Sep 2016 (4th version)	<b>Matsuura Ninjinto Extract Granules</b> Matsuura Yakugyo Co., Ltd.	(61AM)4042 Oct 1986 Oct 1986	6.0	3.0	Hypromellose, Magnesium Stearate, Lactose Hydrate, Dextrin, Corn Starch	Granules	32, Matsuura	A	before or between meals, 2 or 3 times	500g 2g x 84 packets 2g x 300 packets	3.0	3.0	3.0	3.0	3.0
5	Dec 2006 (1st version)	<b>OHSUGI Ninjinto Extract Granules G</b> Ohsugi Pharmaceutical Co., Ltd.	(61AM)4032 Oct 1986 Oct 1986	6.0	2.6	Light Anhydrous Silicic Acid, Lactose Hydrate, Corn Starch, Carmellose Calcium	Granules	SG-32	A	before or between meals, 2 or 3 times	500g 588g (2.0g x 294 packets) 168g (2.0g x 84 packets)	3.0	3.0	3.0	3.0	3.0
6	Revised: Oct 2010 (4th version)	<b>Taikodo Ninjinto Extract Granules</b> Taikoseido Pharmaceutical Co., Ltd.	(61AM)3863 Oct 1986 Nov 1986	7.5	3.56	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-32	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	3.0	3.0	3.0	3.0	3.0
7	Revised: Aug 2017 (6th version)	<b>TEIKOKU Ninjinto Extract Granules</b> Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3681 Oct 1987 Oct 1987	7.5	2.06	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 32	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	3.0	3.0	3.0	3.0	3.0
8	Revised: Oct 2005 (4th version)	<b>Toyo Ninjinto Extract Fine Granules</b> Toyo Yakuko Co., Ltd.	(61AMY)0488 Oct 1987 Feb 1988	4.5	2.4	Corn Starch	Fine granules	TY-088	A	on empty stomach, 3 times	500g 1.5g x 336 packets	3.0	3.0	3.0	3.0	3.0
9	Revised: May 2007 (4th version)	<b>TSUMURA Ninjinto Extract Granules for Ethical Use</b> Tsumura & Co.	(61AM)3276 Oct 1986 Oct 1986	7.5	2.5	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/32	C	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	3.0	3.0	3.0

Table of Indications

A	The following symptoms of those patients with increased urine volume who easily have cold hands and feet: Weak digestive system, gastric atony, diarrhea, vomiting, and gastric pain
B	The following symptoms of those patients with anemia, poor circulation, pressure and heaviness in the stomach, stomachache, tendency to soft stools or diarrhea sometimes accompanied by heaviness of head and vomiting: Chronic diarrhea, gastritis, gastric atony, anemia, autointoxication in physically weak children, and pediatric anorexia
C	The following symptoms of those patients with delicate constitution or those whose weak constitution declined due to disease or fatigue: Acute & chronic enterogastitis, gastric atony, gastric dilation, vomiting of pregnancy (morning sickness), atrophic kidney