

086 Ninjin'yoeito

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio											
												Ginseng	Japanese Angelica Root	Peony Root	Rehmannia Root	Atractylodes Rhizome	Poria Sclerotium	Cinnamon Bark	Astragalus Root	Citrus Unshu Peel	Polypodium Root	Schisandra Fruit	Glycyrrhiza
1	Revised: Apr 2008 (5th version)	<b>Kotaro Ninjin'yoeito Extract Fine Granules</b> Kotaro Pharmaceutical Co., Ltd.	(61AM)3817 Oct 1986 Oct 1986	15.0	9.2	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N108	B	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g)	3.0	4.0	2.0	4.0	4.0	4.0	2.5	1.5	2.0	2.0	1.0	1.0
2	Revised: Jul 2010 (2nd version)	<b>Kracie Ninjin'yoeito Extract Fine Granules</b> Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3510 Jul 2007 Jul 2007	7.5	6.7	JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose, Hydrated Silicon Dioxide	Fine granules	KB-108 3.75g EK-108 2.5g	A	before or between meals, 2 or 3 times	3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g	3.0	4.0	2.0	4.0	4.0	4.0	2.5	1.5	2.0	2.0	1.0	1.0
3	Oct 2005 (1st version)	<b>OHSUGI Ninjin'yoeito Extract Granules G</b> Ohsugi Pharmaceutical Co., Ltd.	(61AM)3871 Oct 1986 Oct 1986	12.0	6.9	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-108	A	before or between meals, 2 or 3 times	500g 1,176g (4.0g x 294 packets) 336g (4.0g x 84 packets)	3.0	4.0	2.0	4.0	4.0	4.0	2.5	1.5	2.0	2.0	1.0	1.0
4	Revised: Mar 2013 (6th version)	<b>TSUMURA Ninjin'yoeito Extract Granules for Ethical Use</b> Tsumura & Co.	(61AM)3305 Oct 1986 Oct 1986	9.0	6.0	JP Magnesium Stearate, JP Lactose Hydrate,	Granules	TSUMURA/108	A	before or between meals, 2 or 3 times	500g 3.0g x 42 packets 3.0g x 189 packets	3.0	4.0	2.0	4.0	4.0	4.0	2.5	1.5	2.0	2.0	1.0	1.0

Table of Indications

A	Declined constitution after recovery from disease, fatigue and malaise, anorexia, perspiration during sleep, cold limbs, and anemia
B	The following symptoms of those patients with ill complexion, thin body, slight fever, chills, persistent cough, severe malaise, anorexia, mental dys-equilibrium, insomnia, night sweating, and tendency to constipation: Enhancement of physical strength after disease or childbirth and delicate constitution