

087 Nyoshinsan

| | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | | | | | | |
|---|--|---|--|----------------------|-------------------------------------|---|-------------|-----------------------------|-------------|--|---|------------------------|----------------|---------------------------|-----------------|---------------|------------------|---------|------|----------------|----------------|-------|
| | | | | | | | | | | | | Japanese Angelica Root | Chidum Rhizome | Aracylodes Lancea Rhizome | Cyperus Rhizome | Cinnamon Bark | Scutellaria Root | Ginseng | Arca | Coptis Rhizome | Saussurea Root | Clove |
| 1 | Revised: Mar 2013 (6th version) | TSUMURA Nyoshinsan Extract Granules for Ethical Use Tsumura & Co. | (61AM)3436 Oct 1986 Oct 1986 | 7.5 | 4.5 | JP Magnesium Stearate, JP Lactose Hydrate | Granules | TSUMURA/67 | A | before or between meals, 2 or 3 times | 500g 2.5g x 42 packets 2.5g x 189 packets | 3.0 | 3.0 | 3.0 | 3.0 | 2.0 | 2.0 | 2.0 | 1.0 | 1.0 | 1.0 | 1.0 |

Table of Indications

| | |
|---|---|
| A | The following symptoms of those patients who have a feeling of hot flush and dizziness: Neurosis before or after childbirth, menstrual irregularity, and automatic imbalance syndrome peculiar to women resembling climacteric disturbance |
|---|---|