

091 Orento

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio								
												Coptis Rhizome	Glycyrrhiza	Processed Ginger	Ginger	Ginseng	Cinnamon Bark	Cinnamon Twig	Jujube	Pinellia Tuber
1	Revised: Jan 2008 (5th version)	Kotaro Orento Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)4036 Oct 1986 Oct 1986	7.5	5.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N120	B	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g) 2.5g x 42 packets (105g)	3.0	3.0	3.0	/	3.0	3.0	/	3.0	6.0
2	Revised: Apr 2011 (5th version)	Taikodo Orento Extract Granules Taikoseido Pharmaceutical Co., Ltd.	(61AM)3857 Oct 1986 Nov 1986	4.5	3.76	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-120	A	before or between meals, 3 times	500g 1.5g x 84 packets 1.5g x 336 packets	3.0	3.0	/	3.0	3.0	3.0	/	3.0	6.0
3	Revised: Jan 2014 (5th version)	Toyo Orento Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AMY)0407 Oct 1986 Jan 1987	6.0	4.0	Corn Starch	Fine granules	TY-009	A	on empty stomach, 3 times	500g 2.0g x 252 packets	3.0	3.0	3.0	/	3.0	/	3.0	3.0	5.0
4	Revised: Mar 2013 (5th version)	TSUMURA Orento Extract Granules for Ethical Use Tsumura & Co.	(61AM)3313 Oct 1986 Oct 1986	7.5	4.0	JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids	Granules	TSUMURA/120	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	3.0	/	3.0	3.0	/	3.0	6.0

Table of Indications

A	The following symptoms of those patients who have heavy stomach feeling, sensation of stomach pressure, and anorexia: Acute gastritis, hangover, and stomatitis
B	The following symptoms of those patients with pressure and heaviness in the stomach, decreased appetite, abdominal pain, nausea, vomiting, oral malodor, and coated tongue accompanied by constipation or diarrhea: Gastrointestinal catarrh, stomatitis, dyspepsia, hyperchlorhydria, and hangover