

091 Orento

| | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | | | | |
|---|--|---|--|----------------------|-------------------------------------|---|---------------|-----------------------------|-------------|---|---|-----------------|-------------|------------------|--------|---------|---------------|---------------|--------|----------------|
| | | | | | | | | | | | | Coptis Rhizome | Glycyrrhiza | Processed Ginger | Ginger | Ginseng | Cinnamon Bark | Cinnamon Twig | Jujube | Pinellia Tuber |
| 1 | Revised: Jan 2008 (5th version) | Kotaro Orento Extract Fine Granules Kotaro Pharmaceutical Co., Ltd. | (61AM)4036 Oct 1986 Oct 1986 | 7.5 | 5.0 | Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate | Fine granules | N120 | B | before or between meals, 2 or 3 times | 500g 2.5g x 189 packets (472.5g) 2.5g x 42 packets (105g) | 3.0 | 3.0 | 3.0 | / | 3.0 | 3.0 | / | 3.0 | 6.0 |
| 2 | Revised: Apr 2011 (5th version) | Taikodo Orento Extract Granules Taikoseido Pharmaceutical Co., Ltd. | (61AM)3857 Oct 1986 Nov 1986 | 4.5 | 3.76 | JP Lactose Hydrate, JP Magnesium Stearate | Granules | TM-120 | A | before or between meals, 3 times | 500g 1.5g x 84 packets 1.5g x 336 packets | 3.0 | 3.0 | / | 3.0 | 3.0 | 3.0 | / | 3.0 | 6.0 |
| 3 | Revised: Jan 2014 (5th version) | Toyo Orento Extract Fine Granules Toyo Yakuko Co., Ltd. | (61AMY)0407 Oct 1986 Jan 1987 | 6.0 | 4.0 | Corn Starch | Fine granules | TY-009 | A | on empty stomach, 3 times | 500g 2.0g x 252 packets | 3.0 | 3.0 | 3.0 | / | 3.0 | / | 3.0 | 3.0 | 5.0 |
| 4 | Revised: Mar 2013 (5th version) | TSUMURA Orento Extract Granules for Ethical Use Tsumura & Co. | (61AM)3313 Oct 1986 Oct 1986 | 7.5 | 4.0 | JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids | Granules | TSUMURA/120 | A | before or between meals, 2 or 3 times | 500g 2.5g x 42 packets 2.5g x 189 packets | 3.0 | 3.0 | 3.0 | / | 3.0 | 3.0 | / | 3.0 | 6.0 |

Table of Indications

| | |
|---|--|
| A | The following symptoms of those patients who have heavy stomach feeling, sensation of stomach pressure, and anorexia: Acute gastritis, hangover, and stomatitis |
| B | The following symptoms of those patients with pressure and heaviness in the stomach, decreased appetite, abdominal pain, nausea, vomiting, oral malodor, and coated tongue accompanied by constipation or diarrhea: Gastrointestinal catarrh, stomatitis, dyspepsia, hyperchlorhydria, and hangover |