

096 Ryokankyomishingeninto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio						
												Poria Sclerotium	Glycyrrhiza	Pinellia Tuber	Processed Ginger	Apricot Kernel	Schisandra Fruit	Asiarum Root
1	Revised: Nov 2007 (5th version)	Kotaro Ryokankyomishingeninto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3820 Oct 1986 Oct 1986	7.5	4.5	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N119	A	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g) 2.5g x 42 packets (105g)	4.0	2.0	4.0	2.0	4.0	3.0	2.0
2	Revised: Mar 2013 (5th version)	TSUMURA Ryokankyomishingeninto Extract Granules for Ethical Use Tsumura & Co.	(62AM)577 Oct 1987 Oct 1987	7.5	4.0	JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids	Granules	TSUMURA/119	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	4.0	2.0	4.0	2.0	4.0	3.0	2.0

Table of Indications

A	The following symptoms of those patients with anemia, oversensitivity to cold, and coughing with profuse sputum accompanied by stridor: Bronchitis, bronchial asthma, cardiac weakness, and kidney disease
---	---