099 Ryutanshakanto

	Date of Making or Revision of Package Insert	Product Name / Manufacturer / Distributor	Approval No. / Date of Listing in the NHI Reimbursement Price / Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio: Japanese Angelica Root	Rehmannia Root	Akebia Stem	Scutellaria Root	Alisma Tuber	Plantago Seed	Japanese Gentian	Gardenia Fruit	Glycyrrhiza	Peony Root	Cnidium Rhizome	Coptis Rhizome	Phellodendron Bark	Forsythia Fruit	Mentha Herb	Glehnia Root and Rhizome
1	Revised: Feb 2018 (6th version)	**JUNKOU Ryutanshakanto FC Extract Fine Granules for Ethical Use** Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00381000 Oct 1986 Oct 1986	9.0	5.6	Corn Starch, Lactose Hydrate	Fine granules	FC 76	A	before or between meals, 2 or 3 times	126g (1.5g x 84 packets)	5.0	5.0	5.0	3.0	3.0	3.0	1.0	1.0	1.0							
2	Revised: Feb 2018 (8th version)	**Kotaro Ryutanshakanto Extract Fine Granules** Kotaro Pharmaceutical Co., Ltd.	(61AM)3963 Oct 1986 Oct 1986	9.0	6.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N76	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	1.5	1.5	1.5	1.5	2.0	1.5	2.0	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
3	Revised: Feb 2018 (5th version)	**SANWA Ryutanshakanto Extract Fine Granules** Sanwa Shoyaku Co., Ltd.	(61AM)3600 Oct 1986 Nov 1986	9.0	5.8	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-14	C	before or between meals, 3 times	500g 3g x 300 packets	5.0	5.0	5.0	3.0	3.0	3.0	1.0	1.0	1.0							
4	Revised: Feb 2018 (6th version)	**Taikodo Ryutanshakanto Extract Fine Granules** Taikoseido Pharmaceutical Co., Ltd.	(62AM)0099 Oct 1987 Oct 1987	7.5	5.0	JP Lactose Hydrate, JP Magnesium Stearate	Powders	TM-76G	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	5.0	5.0	5.0	3.0	3.0	3.0	1.5	1.5	1.5							
5	Revised: Feb 2018 (6th version)	**Taikodo Ryutanshakanto Extract Granules** Taikoseido Pharmaceutical Co., Ltd.	(61AM)4222 Oct 1986 Nov 1986	7.5	5.0	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-76	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	5.0	5.0	5.0	3.0	3.0	3.0	1.5	1.5	1.5							
6	Revised: Feb 2018 (6th version)	**Taikodo Ryutanshakanto Extract Powder** Taikoseido Pharmaceutical Co., Ltd.	(61AM)4823 Oct 1987 Oct 1987	7.5	5.0	JP Lactose Hydrate, JP Magnesium Stearate	Powders	TM-76P	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	5.0	5.0	5.0	3.0	3.0	3.0	1.5	1.5	1.5							
7	Revised: Feb 2018 (7th version)	**Toyo Ryutanshakanto Extract Fine Granules** Toyo Yakuko Co., Ltd.	(61AMY)0386 Oct 1986 Jan 1987	9.0	5.7	Corn Starch	Fine granules	TY-108	A	on empty stomach, 3 times	500g 3.0g x 252 packets	5.0	5.0	5.0	3.0	3.0	3.0	1.5	1.5	1.5							
8	Revised: Feb 2018 (8th version)	**TSUMURA Ryutanshakanto Extract Granules for Ethical Use** Tsumura & Co.	(61AM)1146 Oct 1986 Oct 1986	7.5	5.5	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/76	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	5.0	5.0	5.0	3.0	3.0	3.0	1.0	1.0	1.0							

Table of Indications

A	The following symptoms of those patients with a comparatively strong constitution whose muscles in the lower abdomen are likely to become tense: Micturition pain, feeling of residual urine, turbid urine, and leukorrhea
B	The following symptoms of those patients with comparatively strong constipation: Urethritis, bladder catarrh, vaginitis, genitall eczema, white vaginal discharge, genital itching and pain, and endometritis
C	The following symptoms of those patients with a comparatively strong constitution who have inflammation in the bladder or the urethra or the uterus, micturition pain, and dysuria: Urethritis, cystorrhea, vaginitis, leukorrhea, vulvar eczema, bartholinitis, pruritus vulvae, endometritis, and orchitis