

105 Saikoseikanto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio														
												Bupleurum Root	Japanese Angelica Root	Peony Root	Cnidium Rhizome	Rehmannia Root	Coptis Rhizome	Scutellaria Root	Phellodendron Bark	Gardenia Fruit	Forsythia Fruit	Platyodon Root	Burdock Fruit	Trichosanthes Root	Mentha Herb	Glycyrrhiza
1	Revised: Feb 2018 (7th version)	Kotaro Saikoseikanto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)4035 Oct 1986 Oct 1986	9.0	5.7	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N80	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	2.0	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	
2	Revised: Feb 2018 (7th version)	TEIKOKU Saikoseikanto Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3695 Jul 1988 Jul 1988	9.0	4.48	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 80	A	before meals, 3 times	500g 3.0g x 42 packets 3.0g x 252 packets	2.0	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
3	Revised: Feb 2018 (6th version)	TSUMURA Saikoseikanto Extract Granules for Ethical Use Tsumura & Co.	(61AM)3306 Oct 1986 Oct 1986	7.5	4.75	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/80	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	2.0	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5

Table of Indications

A	The following symptoms in children who tend to be irritable: Neurosis, chronic tonsillitis, and eczema
B	The following symptoms of those patients with delicate constitution and scrofulous constitution: Chronic gastrointestinal disease, anemia, cervical adenitis, hilar lymphadenopathy, tonsillar hypertrophy, neurosis, and eczema