

112 Seishinrenshiin

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio								
												Nelumbo Seed	Ophiopogon Root	Poria Sclerotium	Ginseng	Plantago Seed	Saetularia Root	Astragalus Root	Lycium Bark	Glycyrrhiza
1	Revised: Dec 2016 (3rd version)	JUNKOU Seishinrenshiin FC Extract Fine Granules for Ethical Use Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	1600AMY00417000 Oct 1986 Oct 1986	6.0	4.2	Corn Starch, Lactose Hydrate	Fine granules	FC 111	A	before or between meals, 2 or 3 times	168g (2.0g x 84 packets)	4.0	4.0	4.0	3.0	3.0	3.0	2.0	2.0	2.0
2	Revised: Apr 2006 (5th version)	Toyo Seishinrenshiin Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AMY)0409 Oct 1986 Dec 1986	7.5	5.0	Corn Starch	Fine granules	TY-072	B	on empty stomach, 3 times	500g 2.5g x 252 packets	4.0	4.0	4.0	3.0	3.0	3.0	2.0	2.0	1.5
3	Revised: Mar 2013 (6th version)	TSUMURA Seishinrenshiin Extract Granules for Ethical Use Tsumura & Co.	(61AM)1147 Oct 1986 Oct 1986	7.5	5.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/111	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	4.0	4.0	4.0	3.0	3.0	3.0	2.0	2.0	1.5

Table of Indications

A	The following symptoms of those patients who have general malaise, dry mouth or tongue, and difficulty urinating: Feeling of residual urine, pollakiuria, and micturition pain
B	The following symptoms of those patients who have general malaise, dry mouth or tongue, and difficulty urinating: Feeling of residual urine and micturition pain