

115 Shakanzoto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio									
												Processed Glycyrrhiza	Ginger	Cinnamon Bark	Hemp Fruit	Ajube	Ginseng	Rehmannia Root	Ophiopogon Root	Donkey Glue	Gelatin
1	Revised: Feb 2015 (6th version)	Kotaro Shakanzoto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)4039 Oct 1986 Oct 1986	15.0	9.0*	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N64	A	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g)	3.0	0.8	3.0	3.0	3.0	3.0	6.0	6.0	/	2.0
2	Revised: May 2007 (4th version)	TSUMURA Shakanzoto Extract Granules for Ethical Use Tsumura & Co.	(61AM)3312 Oct 1986 Oct 1986	9.0	7.0	JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/64	B	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 3.0g x 42 packets 3.0g x 189 packets	3.0	1.0	3.0	3.0	3.0	3.0	6.0	6.0	2.0	/

*: extract weight without Gelatin

Table of Indications

A	The following symptoms of those patients with ill complexion, anemia, arrhythmia, severe palpitation and shortness of breath, tendency to constipation, and hot feeling: Cardiac neurosis, cardiac valvulopathy, cough with bloody sputum, and dyspnea in Basedow's disease
B	Palpitation and shortness of breath in patients with a declined constitution who are easily fatigued