

116 Shakuyakukanzobushito

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio		
												Peony Root	Glycyrrhiza	Processed Aconite Root
1	Revised: Mar 2022 (1st version of new form)	SANWA Shakuyakukanzobushito Extract Fine Granules Sanwa Shoyaku Co., Ltd.	16100AMZ03611000 - Nov 1986	4.5	2.6	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-05	A	before or between meals, 3 times	500g [bottle] 1.5g×300 packets	5.0	5.0	1.0
2	Revised: Mar 2017 (5th version)	SANWA Shakuyakukanzobushito Extract Fine Granules Sanwa Shoyaku Co., Ltd. Ohsugi Pharmaceutical Co., Ltd.	(61AM)3611 Oct 1986 Nov 1986	4.5	2.6	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(SG-146)	A	before or between meals, 3 times	500g 441g (1.5g x 294 packets)	5.0	5.0	1.0

Table of Indications

A	The following symptoms of those patients who have oversensitivity to cold, joint pain, myalgia, numbness and extremities that are hard to bend or stretch: Chronic neuralgia, chronic arthritis, articular rheumatism, muscular rheumatism, frozen shoulder, and shoulder stiffness
---	--