

120 Shikunshito

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio							
												Ginseng	Ataractylodes Rhizome	Ataractylodes Rhizome	Poria Sclerotium	Glycyrrhiza	Ginger	Fresh Ginger	Jujube
1	Oct 2005 (1st version)	OHSUGI Shikunshito Extract Tablets Ohsugi Pharmaceutical Co., Ltd.	(61AM)3906 Oct 1986 Oct 1986	5.58 (18 tablets)	3.5	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Carmellose Calcium, Magnesium Stearate, Hypromellose, Titanium Oxide, Food Yellow No.5, Food Blue No.1	Tablets (Film-coated tablets)	SG-75T	A-1	before or between meals, 2 or 3 times	1,764 tablets (6 tablets x 294 packets) 504 tablets (6 tablets x 84 packets)	4.0	4.0	/	4.0	1.5	0.5	/	1.5
2	Revised: Oct 2005 (4th version)	Toyo Shikunshito Extract Fine Granules Toyo Yakuko Co., Ltd.	(62AMY)0396 Jul 1988 Oct 1988	6.0	3.3	Corn Starch	Fine granules	TY-054	A-2	on empty stomach, 3 times	500g 2.0g x 252 packets	4.0	4.0	/	4.0	1.5	/	1.5	1.5
3	Revised: Mar 2013 (5th version)	TSUMURA Shikunshito Extract Granules for Ethical Use Tsumura & Co.	(61AM)1160 Oct 1986 Oct 1986	7.5	2.75	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/75	A-1	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	4.0	/	4.0	4.0	1.0	1.0	/	1.0

Table of Indications

A-1	The following symptoms of those patients who are thin and easily fatigued and have a bad complexion and anorexia: Weak digestive system, chronic gastritis, heavy stomach feeling, vomiting, and diarrhea
A-2	The following symptoms of those patients who are thin and easily fatigued and have a bad complexion and anorexia: Weak digestive system, chronic gastritis, and heavy stomach feeling