

130 Shokenchuto

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio					
												Cinnamon Bark	Ginger	Jujube	Peony Root	Glycyrrhiza	Koi
1	Revised: Feb 2012 (5th version)	Kotaro Shokenchuto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3802 Oct 1986 Oct 1986	27.0	4.5*	Magnesium Stearate, Corn Starch, Pullulan, Magnesium Aluminometasilicate	Fine granules	N99	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g)	4.0	1.0	4.0	6.0	2.0	20.0
2	Revised: Jun 2016 (2nd version)	OHSUGI Shokenchuto Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3885 Oct 1986 Oct 1986	25.2	4.0*	Corn Starch, Sucrose Esters of Fatty Acids	Granules	SG-99	A	before or between meals, 2 or 3 times	500g 1,234.8g (4.2g x 294 packets) 352.8g (4.2g x 84 packets)	4.0	1.0	4.0	6.0	2.0	20.0
3	Revised: May 2007 (4th version)	TSUMURA Shokenchuto Extract Granules for Ethical Use Tsumura & Co.	(61AM)1152 Oct 1986 Oct 1986	15.0	3.75*	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/99	A	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 84 packets 2.5g x 189 packets	4.0	1.0	4.0	6.0	2.0	10.0

*: extract weight without Koi

Table of Indications

A	The following symptoms accompanied by any of delicate constitution and easily fatigued, a poor complexion, abdominal pain, palpitation, hot flushes of the limbs, coldness, pollakiuria, or polyuria: Delicate constitution in childhood, fatigue and malaise, nervousness, chronic gastroenteritis, nocturnal enuresis in children, and night cry
B	The following symptoms of those patients with delicate constitution, fatigability, hot flush, abdominal pain, poor circulation, hot flushes of hands and legs, increased urinary frequency and output: Gastrointestinal disease, pediatric diarrhea or constipation, nervousness, scrofulous constitution, anemia, pollakiuria, pediatric night terrors, and pediatric nocturnal enuresis