

132 Shosai-koto

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio							
												Shojojun Root	Prunella Root	Ginger	Spice Ginger	Scutellaria Root	Jujube	Ginseng	Glycyrrhiza
1	Revised: Apr 2014 (8th version)	Honzo Shosai-koto Extract Granules-M Honzo Pharmaceutical Co., Ltd.	(61AM)4139 Oct 1986 Oct 1986	7.5	4.2	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate	Granules	H09	A	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	7.0	5.0	1.0		3.0	3.0	3.0	2.0
2	Revised: Apr 2014 (9th version)	JPS Shosai-koto Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd.	61AM-4129 Oct 1986 Nov 1986	7.5	4.3	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-09)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	7.0	5.0	1.0		3.0	3.0	3.0	2.0
3	Revised: Dec 2017 (5th version)	JUNKOU Shosai-koto FC Extract Fine Granules for Ethical Use Kohwayakutani Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00362000 Oct 1986 Oct 1986	6.0	3.95	Corn Starch, Lactose Hydrate	Fine granules	FC 09	A	before or between meals, 2 or 3 times	168g (2.0g x 84 packets)	7.0	5.0	1.0		3.0	3.0	3.0	2.0
4	Revised: Apr 2014 (11th version)	Kotaro Shosai-koto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3804 Oct 1986 Oct 1986	7.5	5.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N9	A	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g) 2.5g x 42 packets (105g)	7.0	5.0	1.0		3.0	3.0	3.0	2.0
5	Revised: Apr 2014 (5th version)	Kracie Shosai-koto Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3516 Jul 2007 Jul 2007	6.0	5.4	JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose, Hydrated Silicon Dioxide	Fine granules	KB-9 3.0g EK-9 2.0g	A	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	7.0	5.0	1.0		3.0	3.0	3.0	2.0
6	Revised: Apr 2014 (5th version)	Kracie Shosai-koto Extract Tablets Omimedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00631000 Jul 2007 Jul 2007	5.94 (18 tablets)	4.8	JP Sodium Lauryl Sulfate, JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose, JP Croscarmellose Sodium	Tablets	(EKT-9)	A	before or between meals, 2 or 3 times	252 tablets (3 tablets x 6 packets x 14 sheets) 1,764 tablets (3 tablets x 6 packets x 98 sheets)	7.0	5.0	1.0		3.0	3.0	3.0	2.0
7	Revised: Apr 2016 (8th version)	Matsaura Shosai-koto Extract Granules Matsaura Yakugyo Co., Ltd.	(61AM)4048 Oct 1986 Oct 1986	6.0	viscous extract 7.6*	Hypromellose, Magnesium Stearate, Lactose Hydrate, Dextrin, Wheat Starch	Granules	9, Matsaura	A	before or between meals, 2 or 3 times	500g 2g x 84 packets 2g x 300 packets	7.0	5.0	1.0		3.0	3.0	3.0	2.0
8	Revised: Apr 2014 (3th version)	OHSUGI Shosai-koto Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3876 Oct 1986 Oct 1986	7.5	4.0	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-09	A	before or between meals, 2 or 3 times	500g 735g (2.5g x 294 packets) 210g (2.5g x 84 packets)	7.0	5.0	1.0		3.0	3.0	3.0	2.0
9	Revised: Sep 2016 (4th version)	OHSUGI Shosai-koto Extract T Tablets Takasago Pharmaceutical Industry Co., Ltd. Ohsugi Pharmaceutical Co., Ltd.	(61AM)4190 Oct 1986 Oct 1986	5.94 (18 tablets)	4.0	Microcrystalline Cellulose, Magnesium Aluminometasilicate, Carmellose Calcium, Magnesium Stearate, Hypromellose, Titanium Oxide, Food Yellow No.5 Aluminium Lake, Food Blue No.1 Aluminium Lake, Food Red No.3 Aluminium Lake	Tablets (Film-coated tablets)	(SG-09T)	A	before or between meals, 2 or 3 times	1,764 tablets (6 tablets x 294 packets) 504 tablets (6 tablets x 84 packets)	7.0	5.0	1.0		3.0	3.0	3.0	2.0
10	Revised: Apr 2014 (7th version)	SANWA Shosai-koto Extract Fine Granules Sanwa Shoyaku Co., Ltd.	(61AM)3610 Oct 1986 Nov 1986	7.5	4.6	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-11	A	before or between meals, 3 times	500g 2.5g x 300 packets	6.0	5.0	1.0		3.0	3.0	3.0	2.0
11	Revised: Apr 2014 (8th version)	Taikodo Shosai-koto Extract Granules Taikoseido Pharmaceutical Co., Ltd.	(61AM)3856 Oct 1986 Nov 1986	6.0	4.25	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-9	A	before or between meals, 3 times	500g 2.0g x 84 packets 2.0g x 315 packets	7.0	5.0	1.0		3.0	3.0	3.0	2.0
12	Revised: Aug 2017 (11th version)	TEIKOKU Shosai-koto Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3865 Oct 1987 Oct 1987	7.5	3.45	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 9	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	7.0	5.0	1.0		3.0	3.0	3.0	2.0
13	Revised: Apr 2014 (11th version)	Toyo Shosai-koto Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AMY)0401 Oct 1986 Dec 1986	7.5	5.0	Corn Starch	Fine granules	TY-064	A	on empty stomach, 3 times	500g 2.5g x 252 packets	6.0	5.0		4.0	3.0	3.0	3.0	2.0
14	Revised: Apr 2014 (10th version)	TSUMURA Shosai-koto Extract Granules for Ethical Use Tsumura & Co.	(61AM)3280 Oct 1986 Oct 1986	7.5	4.5	JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids	Granules	TSUMURA 9	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	7.0	5.0	1.0		3.0	3.0	3.0	2.0

*): correspond to 3.8g of dry extract

Table of Indications

A	1. The following symptoms of those patients with moderately strong constitution, right upper abdominal tenderness accompanied by fullness and discomfort, coated tongue, oral cavity discomfort, anorexia, and /or those with slight fever and nausea: Various acute febrile diseases, pneumonia, bronchitis, common cold, lymphadenitis, chronic gastrointestinal disorder, and insufficient postpartum recovery
---	--