

132 Shosaikoto

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio						
												Bupleurum Root	Pinellia Tuber	Ginger	Fresh Ginger	Scutellaria Root	Jujube	Ginseng
1	Revised: Apr 2014 (8th version)	Honzo Shosaikoto Extract Granules-M Honzo Pharmaceutical Co., Ltd.	(61AM)4139 Oct 1986 Oct 1986	7.5	4.2	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminometasilicate, Magnesium Stearate	Granules	H09	A	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	7.0	5.0	1.0	3.0	3.0	3.0	2.0
2	Revised: Apr 2014 (9th version)	JPS Shosaikoto Extract Granules for Ethical Use JPS Pharmaceutical Co., Ltd.	61AM-4129 Oct 1986 Nov 1986	7.5	4.3	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-09)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	7.0	5.0	1.0	3.0	3.0	3.0	2.0
3	Revised: Feb 2022 (1st version of new form)	JUNKOU Shosaikoto FC Extract Fine Granules for Ethical Use Kohwayakutai Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00362000 - Oct 1986	6.0	3.95	Corn Starch, Lactose Hydrate	Fine granules	FC 09	A	before or between meals, 2 or 3 times	168g(2.0g x 84 packets)	7.0	5.0	1.0	3.0	3.0	3.0	2.0
4	Revised: Apr 2014 (11th version)	Kotaro Shosaikoto Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3804 Oct 1986 Oct 1986	7.5	5.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N9	A	before or between meals, 2 or 3 times	500g 2.5g x 189 packets (472.5g) 2.5g x 42 packets (105g)	7.0	5.0	1.0	3.0	3.0	3.0	2.0
5	Revised: Apr 2014 (5th version)	Kracie Shosaikoto Extract Fine Granules Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3516 Jul 2007 Jul 2007	6.0	5.4	JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose, Hydrated Silicon Dioxide	Fine granules	KB-9 3.0g EK-9 2.0g	A	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	7.0	5.0	1.0	3.0	3.0	3.0	2.0
6	Revised: Apr 2014 (5th version)	Kracie Shosaikoto Extract Tablets Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00631000 Jul 2007 Jul 2007	5.94 (18 tablets)	4.8	JP Sodium Lauryl Sulfate, JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose, JP Croscarmellose Sodium	Tablets	(EKT-9)	A	before or between meals, 2 or 3 times	252 tablets (3 tablets x 6 packets x 14 sheets) 1,764 tablets (3 tablets x 6 packets x 98 sheets)	7.0	5.0	1.0	3.0	3.0	3.0	2.0
7	Revised: Apr 2016 (8th version)	Matsuura Shosaikoto Extract Granules Matsuura Yakugyo Co., Ltd.	(61AM)4048 Oct 1986 Oct 1986	6.0	viscous extract 7.6*	Hypromellose, Magnesium Stearate, Lactose Hydrate, Dextrin, Wheat Starch	Granules	9, Matsuura	A	before or between meals, 2 or 3 times	500g 2g x 84 packets 2g x 300 packets	7.0	5.0	1.0	3.0	3.0	3.0	2.0
8	Revised: Feb 2022 (1st version of new form)	OHSUGI Shosaikoto Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	16100AMZ03876000 - Oct 1986	7.5	4.0	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-09	A	before or between meals, 2 or 3 times	500g [bottle] 735g (2.5g x 294 packets) 210g (2.5g x 84 packets)	7.0	5.0	1.0	3.0	3.0	3.0	2.0
9	Revised: Feb 2022 (1st version of new form)	OHSUGI Shosaikoto Extract T Tablets Ohsugi Pharmaceutical Co., Ltd.	16100AMZ04190000 - Oct 1986	5.94 (18 tablets)	4.0	Microcrystalline Cellulose, Magnesium Aluminometasilicate, Carmellose Calcium, Magnesium Stearate, Hypromellose, Titanium Oxide, Food Yellow No.5, Food Blue No.1, Food Red No.3	Tablets (Film-coated tablets)	SG-09T	A	before or between meals, 2 or 3 times	1,764 tablets (6 tablets x 294 packets) 504 tablets (6 tablets x 84 packets)	7.0	5.0	1.0	3.0	3.0	3.0	2.0
10	Revised: Apr 2022 (1st version of new form)	SANWA Shosaikoto Extract Fine Granules Sanwa Shoyaku Co., Ltd.	16100AMZ03610000 - Nov 1986	7.5	4.6	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-11	A	before or between meals, 3 times	500g [bottle] 2.5g x 300 packets	6.0	5.0	1.0	3.0	3.0	3.0	2.0
11	Revised: Apr 2014 (8th version)	Taikodo Shosaikoto Extract Granules Taikoseido Pharmaceutical Co., Ltd.	(61AM)3856 Oct 1986 Nov 1986	6.0	4.25	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-9	A	before or between meals, 3 times	500g 2.0g x 84 packets 2.0g x 315 packets	7.0	5.0	1.0	3.0	3.0	3.0	2.0
12	Revised: Aug 2017 (11th version)	TEIKOKU Shosaikoto Extract Granules Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3865 Oct 1987 Oct 1987	7.5	3.45	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 9	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	7.0	5.0	1.0	3.0	3.0	3.0	2.0
13	Revised: May 2020 (12th version)	Toyo Shosaikoto Extract Fine Granules Toyo Yakuko Co., Ltd.	(61AMY)0401 Oct 1986 Dec 1986	7.5	5.0	Corn Starch	Fine granules	TY-064	A	on empty stomach, 3 times	500g 2.5g x 252 packets	6.0	5.0	4.0	3.0	3.0	3.0	2.0
14	Revised: Apr 2014 (10th version)	TSUMURA Shosaikoto Extract Granules for Ethical Use Tsumura & Co.	(61AM)3280 Oct 1986 Oct 1986	7.5	4.5	JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids	Granules	TSUMURA/9	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	7.0	5.0	1.0	3.0	3.0	3.0	2.0

*: correspond to 3.8g of dry extract

Table of Indications

A	1.The following symptoms of those patients with moderately strong constitution, right upper abdominal tenderness accompanied by fullness and discomfort, coated tongue, oral cavity discomfort, anorexia, and /or those with slight fever and nausea: Various acute febrile diseases, pneumonia, bronchitis, common cold, lymphadenitis, chronic gastrointestinal disorder, and insufficient postpartum recovery
---	---