

136 Tokakujokito

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio					
												Peach Kernel	Cinnamon Bark	Rhubarb	Sodium Sulfate Hydrate	Anhydrous Sodium Sulfate	Glycyrrhiza
1	Revised: Apr 2017 (4th version)	<b>Honzo Tokakujokito Extract Granules-M</b> Honzo Pharmaceutical Co., Ltd.	(61AM)4142 Oct 1986 Oct 1986	7.5	2.3	Lactose Hydrate, Magnesium Aluminometasilicate, Magnesium Stearate	Granules	H61	A-1	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	5.0	4.0	3.0	2.0	1.5	
2	Revised: Apr 2016 (5th version)	<b>JPS Tokakujokito Extract Granules for Ethical Use</b> JPS Pharmaceutical Co., Ltd.	62AM-0477 Oct 1987 Oct 1987	7.5	2.4	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-61)	A-1	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	5.0	4.0	3.0	2.0	1.5	
3	Revised: Feb 2022 (1st version of new form)	<b>JUNKOU Tokakujokito FC Extract Fine Granules for Ethical Use</b> Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00419000 - Oct 1986	6.0	2.3	Corn Starch, Lactose Hydrate	Fine granules	FC 61	A-1	before or between meals, 2 or 3 times	168g (2.0g x 84 pakets)	5.0	4.0	3.0	2.0	1.5	
4	Revised: Oct 2016 (6th version)	<b>Kotaro Tokakujokito Extract Fine Granules</b> Kotaro Pharmaceutical Co., Ltd.	(61AM)3829 Oct 1986 Oct 1986	6.0	3.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N61	B	before or between meals, 2 or 3 times	500g 2.0g x 231 packets (462g) 2.0g x 42 packets (84g)	5.0	4.0	3.0	0.9	1.5	
5	Revised: May 2017 (3rd version)	<b>Kracie Tokakujokito Extract Fine Granules</b> Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3521 Jul 2007 Jul 2007	6.0	2.5	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Light Anhydrous Silicic Acid, JP White Soft Sugar, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-61 3.0g EK-61 2.0g	A-1	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	5.0	4.0	3.0	1.0	1.5	
6	Revised: May 2017 (3rd version)	<b>Kracie Tokakujokito Extract Tablets</b> Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00635000 Jul 2007 Jul 2007	5.94 (18 tablets)	2.2	JP Magnesium Stearate, JP Talc, JP Carmellose Calcium, JP Light Anhydrous Silicic Acid, JP Lactose Hydrate, JP Microcrystalline Cellulose, Magnesium Hydroxide-Aluminium Hydroxide Co-Precipitate	Tablets	(EKT-61)	A-1	before or between meals, 2 or 3 times	252 tablets (3 tablets x 6 packets x 14 sheets) 1,764 tablets (3 tablets x 6 packets x 98 sheets)	5.0	4.0	3.0	1.0	1.5	
7	Revised: Feb 2022 (1st version of new form)	<b>OHSUGI Tokakujokito Extract Granules G</b> Ohsugi Pharmaceutical Co., Ltd.	16100AMZ04810000 - Oct 1987	4.5	2.4	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-61	A-2	before or between meals, 2 or 3 times	500g[bottle] 441g(1.5g x 294 packets) 126g(1.5g x 84 packets)	5.0	4.0	3.0	0.9	1.5	
8	Revised: Aug 2017 (7th version)	<b>TEIKOKU Tokakujokito Extract Granules</b> Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3655 Oct 1987 Oct 1987	7.5	2.38	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 61	A-1	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	5.0	4.0	3.0	2.0	1.5	
9	Revised: May 2007 (4th version)	<b>TSUMURA Tokakujokito Extract Granules for Ethical Use</b> Tsumura & Co.	(61AM)3281 Oct 1986 Oct 1986	7.5	3.0	JP Light Anhydrous Silicic Acid, JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/61	A-1	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	5.0	4.0	3.0	0.9	1.5	

Table of Indications

A-1	The following symptoms of those patients with a comparatively strong constitution and hot flashes who are likely to have constipation: Menstrual irregularity, dysmenorrhea, anxiety during menstruation or following childbirth, low back pain, constipation, accessory symptoms associated with hypertension (headache, dizziness, and shoulder stiffness)
A-2	The following symptoms of those patients with a comparatively strong constitution and hot flashes who are likely to have constipation: Menstrual irregularity, dysmenorrhea, anxiety during menstruation or following childbirth, low back pain, constipation, accessory symptoms associated with hypertension (headache, shoulder stiffness and dizziness)
B	The following symptoms of those patients with headache, tendency to hot flush, tenderness and fecal impaction in the left lower abdomen, cold limbs and lower back, and decreased urinary output: Chronic constipation, hypertension, arteriosclerosis, lower back pain, hemorrhoids, various disorders due to maenstrual irregularity, climacteric disturbance, acne, blot, eczema, white vaginal discharge, and sciatica