

139 Tokishakuyakusan

No.	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio						
												Japanese Angelica Root	Chilium Rhizome	Penny Root	Poria Sclerotium	Anacardios Rhizome	Anacardios/Lancea Rhizome	Aisatsu Tuber
1	Revised: Oct 2011 (4th version)	<b>Honzo Tokishakuyakusanryo Extract Granules-M</b> Honzo Pharmaceutical Co., Ltd.	(61AM)4161 Oct 1986 Oct 1986	7.5	4.5	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Aluminummetasilicate, Magnesium Stearate	Granules	H23	A	before or between meals, 3 times	500g 2.5g x 294 packets 2.5g x 42 packets	3.0	3.0	4.0	4.0	4.0	4.0	
2	Revised: Sep 2012 (5th version)	<b>JPS Tokishakuyakusanryo Extract Granules for Ethical Use</b> JPS Pharmaceutical Co., Ltd.	61AM-4766 Oct 1987 Oct 1987	7.5	4.6	Magnesium Stearate, Sucrose Esters of Fatty Acids, Lactose Hydrate	Granules	(J-23)	A	before or between meals, 2 or 3 times	500g 750g (2.5g x 300 packets) 105g (2.5g x 42 packets)	3.0	3.0	4.0	4.0	4.0	4.0	
3	Revised: Oct 2016 (3rd version)	<b>JUNKOU Tokishakuyakusanryo FC Extract Fine Granules for Ethical Use</b> Kohwayakutuu Inc. Ohsugi Pharmaceutical Co., Ltd.	16100AMY00364000 Oct 1986 Oct 1986	6.0	3.65	Corn Starch, Lactose Hydrate	Fine granules	FC 23	A	before or between meals, 2 or 3 times	168g (2.0g x 84 packets)	3.0	3.0	6.0	4.0	4.0	5.0	
4	Revised: Nov 2010 (6th version)	<b>Kotaro Tokishakuyakusanryo Extract Fine Granules</b> Kotaro Pharmaceutical Co., Ltd.	(61AM)3830 Oct 1986 Oct 1986	9.0	5.5	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminummetasilicate	Fine granules	N23	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	3.0	3.0	4.0	4.0	4.0	4.0	
5	Revised: Oct 2013 (4th version)	<b>Kracie Tokishakuyakusanryo Extract Fine Granules</b> Kracie Pharma, Ltd. Kracie Pharmaceutical, Ltd.	(61AM)3518 Jul 2007 Jul 2007	6.0	5.0	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-23 3.0g EK-23 2.0g	A	before or between meals, 2 or 3 times	3.0g x 28 packets 3.0g x 168 packets 2.0g x 42 packets 2.0g x 294 packets 500g	3.0	3.0	6.0	4.0	4.0	4.0	
6	Revised: Apr 2016 (4rd version)	<b>Matsuura Tokishakuyakusanryo Extract Granules</b> Matsuura Yakugyo Co., Ltd.	(61AM)4173 Oct 1986 Oct 1986	7.5	viscous extract 8.4*	Hypromellose, Magnesium Stearate, Lactose Hydrate, Dextrin, Corn Starch	Granules	23, Matsuura	A	before or between meals, 2 or 3 times	500g 2.5g x 84 packets 2.5g x 300 packets	3.0	3.0	4.0	4.0	4.0	4.0	
7	Oct 2005 (1st version)	<b>OHSUGI Tokishakuyakusanryo Extract Granules G</b> Ohsugi Pharmaceutical Co., Ltd.	(61AM)4811 Oct 1987 Oct 1987	7.5	4.2	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-23	A	before or between meals, 2 or 3 times	500g 735g (2.5g x 294 packets) 210g (2.5g x 84 packets)	3.0	3.0	4.0	4.0	4.0	4.0	
8	Jan 2019 (1st version)	<b>OHSUGI Tokishakuyakusanryo Extract T Tablets</b> Ohsugi Pharmaceutical Co., Ltd.	(61AM)4786 Oct 1987 Oct 1987	5.94 (18 tablets)	4.2	Microcrystalline Cellulose, Magnesium Aluminummetasilicate, Carmellose Calcium, Magnesium Stearate, Hypromellose, Titanium Oxide, Food Yellow No.5 Aluminium Lake, Food Blue No.1 Aluminium Lake, Food Red No.3 Aluminium Lake	Tablets (Film-coated tablets)	SG-23T	A	before or between meals, 2 or 3 times	1,764 tablets (6 tablets x 294 packets) 504 tablets (6 tablets x 84 packets)	3.0	3.0	4.0	4.0	4.0	4.0	
9	Revised: Apr 2013 (4th version)	<b>SANWA Tokishakuyakusanryo Extract Fine Granules</b> Sanwa Shoyaku Co., Ltd.	61AM3618 Oct 1986 Nov 1986	7.5	4.8	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-22	C	before or between meals, 3 times	500g 2.5g x 300 packets	3.0	3.0	4.0	4.0	4.0	4.0	
10	Revised: Nov 2011 (6th version)	<b>Taikodo Tokishakuyakusanryo Extract Granules</b> Taikoseido Pharmaceutical Co., Ltd.	(61AM)4408 Oct 1987 Oct 1987	7.5	5.15	JP Lactose Hydrate, JP Magnesium Stearate	Granules	TM-23	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	3.0	3.0	6.0	4.0	4.0	4.0	
11	Revised: Nov 2011 (4th version)	<b>Taikodo Tokishakuyakusanryo Extract Powder</b> Taikoseido Pharmaceutical Co., Ltd.	(61AM)4407 Oct 1987 Oct 1987	7.5	5.15	JP Lactose Hydrate, JP Magnesium Stearate	Powders	TM-23P	A	before or between meals, 3 times	500g 2.5g x 84 packets 2.5g x 252 packets	3.0	3.0	6.0	4.0	4.0	4.0	
12	Revised: Aug 2017 (7th version)	<b>TEIKOKU Tokishakuyakusan Extract Granules</b> Teikoku Kampo Seiyaku Co., Ltd. Teikoku Seiyaku Co., Ltd.	(61AM)3704 Oct 1987 Oct 1987	7.5	3.2	Lactose Hydrate, Microcrystalline Cellulose, Magnesium Stearate	Granules	TEIKOKU 23	A	before meals, 3 times	500g 2.5g x 42 packets 2.5g x 252 packets	3.0	3.0	4.0	4.0	4.0	4.0	
13	Revised: May 2011 (6th version)	<b>Toyo Tokishakuyakusanryo Extract Fine Granules</b> Toyo Yakuko Co., Ltd.	(61AMY)0458 Oct 1987 Feb 1988	7.5	4.2	Corn Starch	Fine granules	TY-085	A	on empty stomach, 3 times	500g 2.5g x 252 packets	3.0	3.0	4.0	4.0	4.0	4.0	
14	Revised: Oct 2014 (6th version)	<b>TSUMURA Tokishakuyakusan Extract Granules for Ethical Use</b> Tsumura & Co.	(61AM)1166 Oct 1986 Oct 1986	7.5	4.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/23	D	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	4.0	4.0	4.0	4.0	

\*: correspond to 4.2g of dry extract

Table of Indications

A	The following symptoms of those patients with comparatively weak constitution, slight oversensitivity to cold, and slight anemia who are easily fatigued and sometimes have lower abdominal pain, dull headache, dizziness, shoulder stiffness, buzzing in the ears, palpitation, etc.: Menstrual irregularity, abnormal menstruation, menses painful, climacteric disturbance, disorders fatigue before/after childbearing or abortion (anemia, fatigue and malaise, dizziness, edema), dizziness, dull headache, shoulder stiffness, low back pain, cold feeling in the lower limbs and waist, chilblain, edema, and spots
B	The following symptoms of those patients with anemia, poor circulation, weak stomach, dark circles around the eyes, fatigability, heaviness of head, dizziness, shoulder stiffness, palpitation, increased urinary frequency, decreased urinary output, thirst or lower abdominal tenderness or pain due to poor circulation or tendency to frostbite: Cardiac debility, kidney disease, anemia, anemia before or after childbirth or after abortion, hemorrhoids, anal prolapse, morning sickness, menstrual irregularity, menstrual pain, climacteric neurosis, acne, blot, and abnormal blood pressure
C	The following symptoms of those patients with anemia, oversensitivity to cold, a bad complexion, unidentified complaints such as dull headache, dizziness, shoulder stiffness, palpitation, cold feeling in the lower limbs and waist, increased urinary frequency and decreased urine volume, lower abdominal pain: Anemia, oversensitivity to cold, climacteric disturbance, infertility, habitual abortion, nephropathy of pregnancy, nephrosis, menstrual irregularity, endometritis, abnormal blood pressure, anal prolapse, and acne vulgaris
D	The following symptoms of those patients who have generally weak muscles and are easily fatigued and whose waist and lower limbs are susceptible to cold: Anemia, malaise, climacteric disturbance (dull headache, headache, dizziness, shoulder stiffness, etc.), menstrual irregularity, dysmenorrhea, infertility, palpitation pounding, chronic nephritis, diseases during pregnancy (edema, habitual abortion, hemorrhoids, abdominal pain), beriberi, hemiplegia and valvular diseases of the heart