

140 Tokishakuyakusankabushi

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio						
												Japanese Angelica Root	Cnidium Rhizome	Peony Root	Poria Sclerotium	Atractylodes Rhizome	Alisma Tuber	Processed Aconite Root
1	Revised: Feb 2016 (4rd version)	<b>SANWA Tokishakuyakusankabushi Extract Fine Granules</b> Sanwa Shoyaku Co., Ltd.	(61AM)3613 Oct 1986 Nov 1986	9.0	5.9	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	S-29	A	before or between meals, 3 times	500g 3g x 300 packets	3.0	3.0	6.0	4.5	4.5	3.5	1.0
2	Revised: Aug 2009 (4th version)	<b>SANWA Tokishakuyakusankabushi Extract Fine Granules</b> Sanwa Shoyaku Co., Ltd. Ohsugi Pharmaceutical Co., Ltd.	(61AM)3613 Oct 1986 Nov 1986	9.0	5.9	Lactose Hydrate, Corn Starch, Microcrystalline Cellulose, Partly Pregelatinized Starch, Light Anhydrous Silicic Acid	Fine granules	(SG-143)	A	before or between meals, 3 times	500g 882g (3g x 294 packets)	3.0	3.0	6.0	4.5	4.5	3.5	1.0

**Table of Indications**

A	The following symptoms of those patients who have a bad complexion, anemia, getting cold easily in the lower limbs and waist, headache, dull headache, increased urinary frequency, occasional dizziness, shoulder stiffness, tinnitus and palpitation: Oversensitivity to cold in women, menorrhagia, neuralgia, chronic nephritis, climacteric disturbance, disorders during pregnancy (edema, prevention of habitual abortion, hemorrhoids, abdominal pain), and slow recovery from the postpartum
---	--