

141 Tokishigyakukagoshuyushokyo

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio								
												Japanese Angelica Root	Cinnamon Bark	Peony Root	Akebia Stem	Asiasarum Root	Glycyrrhiza	Jujube	Euodia Fruit	Ginger
1	Revised: Jun 2009 (6th version)	Kotaro Tokishigyakukagoshuyushokyo Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)4131 Oct 1986 Oct 1986	9.0	6.0	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N38	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	3.0	3.0	3.0	3.0	2.0	2.0	5.0	2.0	1.0
2	Revised: Jul 2010 (3rd version)	Kracie Tokishigyakukagoshuyushokyo Extract Fine Granules Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00620000 Jul 2007 Jul 2007	7.5	4.2	JP Magnesium Stearate, JP Light Anhydrous Silicic Acid, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-38 3.75g EK-38 2.5g	A	before or between meals, 2 or 3 times	3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g	3.0	3.0	3.0	3.0	2.0	2.0	5.0	2.0	1.0
3	Revised: Jun 2009 (2nd version)	OHSUGI Tokishigyakukagoshuyushokyo Extract Granules G Ohsugi Pharmaceutical Co., Ltd.	(61AM)3874 Oct 1986 Oct 1986	9.0	4.6	Lactose Hydrate, Corn Starch, Magnesium Stearate	Granules	SG-38	A	before or between meals, 2 or 3 times	500g 882g (3.0g x 294 packets) 252g (3.0g x 84 packets)	3.0	3.0	3.0	3.0	2.0	2.0	5.0	2.0	1.0
4	Revised: Jun 2009 (5th version)	TSUMURA Tokishigyakukagoshuyushokyo Extract Granules for Ethical Use Tsumura & Co.	(61AM)1125 Oct 1986 Oct 1986	7.5	4.0	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/38	A	before or between meals, 2 or 3 times	500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	3.0	3.0	2.0	2.0	5.0	2.0	1.0

Table of Indications

A	The following symptoms of those patients who have cold hands and feet and are likely to have pain in the lower limbs or lower abdomen when their lower limbs are cold: Chilblain, headache, lower abdominal pain, and low back pain
B	The following symptoms of those patients with anemia, poor circulation, headache, pressure and heaviness in the stomach, low back pain, lower abdominal pain, and tendency to frostbite: Frostbite, chronic headache, sciatica, and lower abdominal pain in women