

143 Tsudosan

| | Date of Making or Revision of Package Insert | Product Name Manufacturer Distributor | Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan | Dose for One Day (g) | Extract Content in One Day Dose (g) | Inactive Ingredients | Dosage Form | ID Code (Item Number, etc.) | Indications | Daily Dosage and Administration (Adult) | Packaging | Component Ratio | | | | | | | | | |
|---|--|---|--|----------------------|-------------------------------------|---|---------------|-----------------------------|-------------|---|--|------------------------|---------|--------------------------|-----------------|---------------|--------------------|-------------|-----------|-------------|------------|
| | | | | | | | | | | | | Japanese Angelica Root | Rhubarb | Anhydrous Sodium Sulfate | Immature Orange | Magnolia Bark | Citrus Unshiu Peel | Akebia Stem | Safflower | Sappan Wood | Glycythiza |
| 1 | Revised: Oct 2016 (8th version) | Kotaro Tsudosan Extract Fine Granules Kotaro Pharmaceutical Co., Ltd. | (61AM)3828 Oct 1986 Oct 1986 | 12.0 | 6.5 | Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate | Fine granules | N105 | A-1 | before or between meals, 2 or 3 times | 500g 4.0g x 147 packets (588g) 4.0g x 42 packets (168g) | 3.0 | 3.0 | 1.8 | 3.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 |
| 2 | Revised: Mar 2020 (6th version) | Taikodo Tsudosan Extract Granules Taikoseido Pharmaceutical Co., Ltd. | (61AM)3858 Oct 1986 Nov 1986 | 7.5 | 5.0 | JP Lactose Hydrate, JP Magnesium Stearate | Granules | TM-105 | A-2 | before or between meals, 3 times | 500g 2.5g x 84 packets 2.5g x 252 packets | 3.0 | 3.0 | 2.0 | 3.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 |
| 3 | Revised: Jun 2009 (5th version) | TSUMURA Tsudosan Extract Granules for Ethical Use Tsumura & Co. | (61AM)1145 Oct 1986 Oct 1986 | 7.5 | 4.5 | JP Magnesium Stearate, JP Lactose Hydrate | Granules | TSUMURA/105 | A-1 | before or between meals, 2 or 3 times | 500g 5kg (500g x 10) 2.5g x 42 packets 2.5g x 189 packets | 3.0 | 3.0 | 1.8 | 3.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 | 2.0 |

Table of Indications

| | |
|-----|---|
| A-1 | The following symptoms of those patients with a comparatively strong constitution who have tenderness in the lower abdomen and are likely to have constipation: Menstrual irregularity, menalgia, climacteric disturbance, low back pain, constipation, bruise (contusion) , and symptoms associated with hypertension (headache, dizziness, and shoulder stiffness) |
| A-2 | The following symptoms of those patients with a comparatively strong constitution who have tenderness in the lower abdomen and are likely to have constipation: Menstrual irregularity, menalgia, climacteric disturbance, constipation, bruise (contusion) , and symptoms associated with hypertension (headache, dizziness, and shoulder stiffness) |