

148 Yokukansankachimpihange

	Date of Making or Revision of Package Insert	Product Name Manufacturer Distributor	Approval No. Date of Listing in the NHI Reimbursement Price Date of Initial Marketing in Japan	Dose for One Day (g)	Extract Content in One Day Dose (g)	Inactive Ingredients	Dosage Form	ID Code (Item Number, etc.)	Indications	Daily Dosage and Administration (Adult)	Packaging	Component Ratio									
												Japanese Angelica Root	Uncaria Hook	Cnidium Rhizome	Atractylodes Rhizome	Atractylodes Lancea Rhizome	Poria Sclerotium	Bupleurum Root	Glycyrrhiza	Citrus Unshiu Peel	Pinellia Tuber
1	Revised: Apr 2008 (6th version)	Kotaro Yokukansankachimpihange Extract Fine Granules Kotaro Pharmaceutical Co., Ltd.	(61AM)3840 Oct 1986 Oct 1986	9.0	6.1	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate	Fine granules	N83	B	before or between meals, 2 or 3 times	500g 3.0g x 168 packets (504g) 3.0g x 42 packets (126g)	3.0	3.0	3.0	4.0	/	4.0	2.0	1.5	3.0	5.0
2	Revised: Jul 2010 (2nd version)	Kracie Yokukansankachimpihange Extract Fine Granules Ominedo Pharmaceutical Industry Co., Ltd. Kracie Pharmaceutical, Ltd.	21900AMX00648000 Jul 2007 Jul 2007	7.5	5.0	JP Magnesium Stearate, JP Microcrystalline Cellulose, JP Lactose Hydrate, Hydrated Silicon Dioxide	Fine granules	KB-83 3.75g EK-83 2.5g	A	before or between meals, 2 or 3 times	3.75g x 28 packets 3.75g x 168 packets 2.5g x 42 packets 2.5g x 294 packets 500g	3.0	3.0	3.0	4.0	/	4.0	2.0	1.5	3.0	5.0
3	Revised: Mar 2013 (5th version)	TSUMURA Yokukansankachimpihange Extract Granules for Ethical Use Tsumura & Co.	(61AM)1129 Oct 1986 Oct 1986	7.5	4.5	JP Magnesium Stearate, JP Lactose Hydrate	Granules	TSUMURA/83	A	before or between meals, 2 or 3 times	500g 2.5g x 42 packets 2.5g x 189 packets	3.0	3.0	3.0	/	4.0	4.0	2.0	1.5	3.0	5.0

Table of Indications

A	The following symptoms of those patients with a delicate constitution and nervousness: Neurosis, insomnia, night cry in children, and peevishness in children
B	Neurosis, climacteric neurosis, insomnia, neurological symptoms due to hypertension or arteriosclerosis, and pediatric night terrors