

Indications

Following symptoms in persons with moderate physical strength who suffer from chronically persistent conditions such as headache, dizziness and stiff shoulders:

Chronic headache, neurosis and tendency to hypertension.

! Caution for use

To be avoided

Don't give this medicine to children under 3 months of age.

Who should seek consultation

- The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - Persons undergoing medical treatment from a physician.
 - Pregnant women or women suspected of being pregnant.
 - Persons with weak stomach and intestine, and sensitiveness to cold.
 - Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
- The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Loss of appetite, gastric distress

- Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.

Components

Chotosan (釣藤散) is extract from mixture of the following crude drugs;

Uncaria Hook (釣藤鈎), Citrus Peel (橘皮) (or Citrus Unshiu Peel (陳皮)), Pinellia Tuber (半夏),
Ophiopogon Root (麦門冬), Poria Sclerotium (茯苓), Ginseng (人參), Saposhnikovia Root and Rhizome (防風),
Chrysanthemum Flower (菊花), Glycyrrhiza (甘草), Ginger (生姜), Gypsum (石膏).



*If you have any questions, please consult to a physician, pharmacist, or registered distributor .

Supervised by Ichiro Arai Ph.D., Professor, Division of Kampo Pharmaceutical Sciences, Nihon Pharmaceutical University



JAPAN KAMPO MEDICINES MANUFACTURERS ASSOCIATION